## BRITISH JOURNAL OF RHEUMATOLOGY – GENERAL INDEX – VOLUME 38

(S1 and S2 = Supplements; E = Editorial; L = Letter to the Editor)

AARONS L see Herrick AL

ABINADER A, HANLY AJ, LOZADA CJ. Catastrophic antiphospholipid syndrome associated with anti-beta-2-glycoprotein I IgA, (L) 84

ACCARDO S see Seriolo B

Acroparasthesia-a typical finding in vitamin D deficiency, (L) 482 Acupuncture

as an adjunctive therapy in osteoarthritis of the knee. A randomized controlled study, 346

effect of on patients with rheumatoid arthritis: a randomized, placebocontrolled cross-over study, 864

for neck pain: a systematic review of randomized controlled trials, 143 Adamantiades-Behçet disease see Behçet's disease

ADAMS MA see Mannion AF

ADLER CP see Jacki SH

Adult Still's disease

high serum level of macrophage-colony stimulating factor (M-CSF) in. (L) 477

with nephrotic syndrome at presentation, (L) 283

and pregnancy, intravenous immunoglobulins for, (L) 1024

AGGARWAL P see Handa R

Ано K see Jäntti J; Myllykangas-Luosujärvi R

AIGIANI MG see Ruperto N AKAHOSHI M see Nakashima H

AKOĞLU T see Hamuryudan V AKSU K, KABASAKAL Y, SAYINER A, KESER G, OKSEL F, BILGIÇ A, GÜMÜŞDIŞ G, Doğanavşargıl E. Prevalences of hepatitis A, B, C and E viruses in Behçet's disease, 1279

AKTUGLU Y see Basaranoglu M

AL-ADSANI A, NIAZY MN, MOHD M. Ewing's sarcoma of the ilium mimicking sacroiliitis, (L) 792

AL-ALI N see Al-Awadhi AM

AL-AWADHI AM, OLUSI SO, AL-ZAID NS, PRABHA K, AL-ALI N, AL-JARALLAH KH, GEORGE S. Prevalence of hypothyroidism among Arabs with rheumatoid arthritis. (L) 472

AL-JARALLAH KH see Al-Awadhi AM

AL-ZAID NS see Al-Awadhi AM

ALAADDINE N see Pelletier J-P

ALBERS JMC, KUPER HH, VAN RIEL PLCM, PREVOO MLL, VAN 'T HOF MA, VAN GESTEL AM, SEVERENS JL. Socio-economic consequences of rheumatoid arthritis in the first years of the disease, 423

ALBERS JMC see also Barrera P

Alcohol-related deaths, rates of from accidents and violence in subjects with ankylosing spondylitis and rheumatoid arthritis, (L) 91

Alessio M see Ruperto N

ALEXANDERSON H, STENSTRÖM CH, LUNDBERG I. Safety of a home exercise programme in patients with polymyositis and dermatomyositis: a pilot study, 608

ALFONSO F see Fernández-Gutiérrez B

Alfuzosin is not associated with dermatomyositis, (L) 579

ALLANORE Y see Claudepierre P

ALLDERSEA JE see Layton MA

Allopurinol hypersensitivity

familial, with subsequent desensitization, (L) 1301

oral desensitization in patients with chronic tophaceous gout and, (L) 85

ALVES J see Ames PRJ

AMENGUAL O see Atsumi T

AMES PRJ, ALVES J, MURAT I, ISENBERG DA, NOUROOZ-ZAHED J. Oxidative stress in systemic lupus erythematosus and allied conditions with vascular involvement, 529

Aminobisphosphonate ibandronate, pro-inflammatory effects in vitro and in vivo. 984

AMOR B see E! Maghraoui A

Amos N see Richards PJ

Amos RS see Squirrell DM

Amyloidosis in a nationwide series of 1666 subjects with rheumatoid arthritis who died during 1989 in Finland, 499

ANDERS H-J, SCHATTENKIRCHNER M. Destructive joint lesions and bursitis in idiopathic hypereosinophilic syndrome, (L) 185

ANDERSON J see Hamilton J

Angio-neurotic oedema associated with methotrexate treatment in rheumatoid arthritis, 908

Angiogenesis and arthritis, 103

Ankle arthropathy, predominant, in hereditary haemochromatosis, (L) 378 Ankle joint complex. A preliminary study determining the feasibility of electromagnetic tracking for kinetics at the, 1260

Ankylosing spondylitis

biochemical analysis of posture in patients with spinal kyphosis due to: a pilot study, 441

cutaneous necrotizing vasculitis, erythema nodosum and, (L) 377

defining disease activity in: is a combination of variables (Bath Ankylosing Spondylitis Disease Activity Index) an appropriate

increased antibody responses to Klebsiella serotypes K26, K36 and K50 in. (L) 481

relationship between urinary pyridinium cross-links, disease activity and disease subsets of, 21

what is the optimum duration of a clinical study? A one year versus a six weeks non-steroidal anti-inflammatory drug trial, 235

Anticardiolipin antibodies

free protein S levels and thrombosis: a survey in a selected population of rheumatoid arthritis patients, 675

relationships between serum 17-\(\beta\)-oestradiol and, in female patients with rheumatoid arthritis, (L) 1159

Weber-Christian disease associated with lupus anticoagulant and, (L)

Antinuclear antibodies and lupus: causes and consequences, (E) 691 Antiphospholipid syndrome

animal models, 59!

catastrophic, associated with anti-beta-2-glycoprotein I IgA, (L) 84 microscopic polyangiitis associated with, (L) 478

primary, correlation between \(\beta^2\)-glycoprotein I valine leucine 247 polymorphism and anti-β2-glycoprotein I antibodies, 721

splenectomy for refractory thrombocytopenia in, 848

APAYDIN H see Ertan S APAYDIN S see Basaranoglu M

Apocynin, a plant-derived, cartilage-saving drug, might be useful in the

treatment of rheumatoid arthritis, 1088 Apoptosis

cell, in synovial membranes in psoriatic, reactive and rheumatoid arthritis. A comparative quantitative morphometric study of, 431 role of in systemic lupus erythmatosus, 1177

APPELBOOM T see Steinfeld SD

ARAL O see Gül A

ARAÚJO O see Garcia-Pardo G

ARENDZEN JH see Winters JC

ARINOBU Y see Nakashima H

ARMADANS-GIL L see Marsal S ARMSTRONG R, RASKER H, DEQUEKER J. An insight into rheumatology resources available on the World Wide Web (http://www.ilar.org), (1.) 1028

AROMAA A see Mäkelä M

ARPACIOĞLU O see Özgül A

Arthritis

acute microcrystalline, seasonal variation in the onset of, 1003

angiogenesis and, 103

antigen-induced, liposomal clodronate eliminates synovial macrophages, reduces inflammation and ameliorates joint destruction

bibliometric methods for the evaluation of research, 13

crystal, management of, (E) 912

detection of mycobacteria in joint samples from patients using a genusspecific polymerase chain reaction and sequence analysis, 547

depressed proliferative responses by peripheral blood mononuclear cells from patients to mycobacterial heat shock protein 60, 631 immunohistological analysis of synovial tissue for differential diagnosis in, 1074

early inflammatory, collagenase, cathepsin B and cathepsin L gene expression in the synovial membrane, 34

effects of interferon beta treatment on, 362

eosinophilic 1188

glomerular and tubular proteinuria as markers of nephropathy in, 28 inflammatory, granisetron (Kytril) suppresses methotrexate-induced nausea and vomiting and is superior to prochlorperazine (Stemetil), 280

and interleukin 10, (E) 293

peripheral benzodiazepine receptor ligand PK 11195 inhibits in the MRL-1pr mouse model, 1068

psoriatic see Psoriatic arthritis

quantitation of interferon gamma- and interleukin-4-producing T cells in synovial fluid and peripheral blood of patients, 214 rabbit antigen-induced, effects of soluble interleukin-1 type II receptor

on: clinical, biochemical and histological asses

Chlamydia pneumoniae as a triggering infection, 411

comparative quantitative morphometric study of cell apoptosis in synovial membranes in psoriatic, and rheumatoid arthritis, 431 rheumatoid see Rheumatoid arthritis

septic

characteristics of in human immunodeficiency virus-infected haemophiliacs versus other risk groups, 139

melioidosis presenting as, in Bengali men in East London,(L) 1029 seronegative, cutaneous polyarteritis nodosa with (L) 1161 Type IX collagen immunoreactive peptides in synovial fluids from

patients, 338 ARTHUR V see Devlin J

Articular hypermobility in Icelandic 12-year-olds, 1014

ARVIDSON N-G see Torsteinsdéttir I

ASAHARA H see Maeda T Asai K see Muro H

ASAMOTO K see Wada M

ASH G, DICKENS CM, CREED FH, JAYSON MIV, TOMENSON B. The effects of dothiepin on subjects with rheumatoid arthritis and depression, 959

ASHTON B see Middleton J ASLANTAS AB see Gül A

Assimadi JK see Mijiyawa M

ATKINS D, CRAWFORD F, EDWARDS J, LAMBERT M. A systematic review of treatments for the painful heel, 968

ATKINS D see also Crawford F

ATSUMI T, TSUTSUMI A, AMENGUAL O, KHAMASHTA MA, HUGHES GRV, Міуоsні Y, Існікаwa K, Коїке T. Correlation between β2-glycoprotein I valine leucine<sup>247</sup> polymorphism and anti-β2-glycoprotein I antibodies in patients with primary antiphospholipid syndrome, 721

ATTOH-MENSAH K see Mijiyawa M ATTUIL G see Seror P

AUBARD Y see Liozon E

AUDRING H see Riemekasten G AUGUET JL see Garcia-Pardo G

Autoimmune congenital heart block, enhanced frequency in female offspring, (L) 380

Autoimmune myelofibrosis and Sjögren's syndrome: an uncommon association, (L) 370

Autologous stem cell transplantation see Stem cell transplantation

Azathioprine, identification of thiopurine methyltransferase (TPMT) polymorphisms cannot predict myelosuppression in systemic lupus erythematosus patients taking, 640

AZIZ W, YATES DB. Hypertrophic osteoarthropathy associated with bacterial endocarditis, (L) 375

BABINET P see Généreau T

Back muscle strength, maximal, prediction from indices of body mass and fat-free body mass, 652

BADER DL see Harms MC

BAGG J see Jackson MS

BAJOCCHI G see Trotta F

BAKKALOGLU A see Yetgin S BAKKER B see Vossen JM

BALACCO GABRIELI C see Giorgi D

BALINT G see Kirwan JR

BALSA A see Villaverde V

BANHAM S see Hamilton J

BARDARE M see Ruperto N

BARETA J see Berman BM BARKER S see Woodburn J

BARNES CG, YAZICI H. Behçet's syndrome, (E) 1171

BARNES L see Flanagan N

BARRERA P, RADSTAKE TRDJ, ALBERS JMC, VAN RIEL PLCM, VAN DE PUTTE LBA. Familial aggregation of rheumatoid arthritis in The Netherlands: a cross-sectional hospital-based survey, 415

BARRETT JH see Donn RP

BARTHELMES H see Riemekasten G

BARZILAI A, VARON D, MARTINOWITZ U, HEIM M, SCHULMAN S. Characteristics of septic arthritis in human immunodeficiency virusinfected haemophiliacs versus other risk groups, 139

BASARANOGLU M, MERT A, TABAK F, APAYDIN S, AKTUGLU Y, OZDOGAN H. A case of familial Mediterranean fever and polyarteritis nodosa complicated by spontaneous bilateral perirenal and subcapsular splenic haemorrhage, (L) 794

BASLER H-D see Dieppe P

BATLLE E see Fernández-Gutiérrez B

BATTAGLIA E see Naughton MA

BEAUCHARD C see Rolf C

Behçet's disease

childhood, clinical features and comparison with adult-onset disease,

evaluation of a new instrument to measure clinical activity, 728

evaluation of the Turkish translation of a disease activity form for, 734 false aneurysm of the internal carotid artery in: successful combined endovascular treatment with stent and coils, (L) 576

lung function in. (L) 1018

organizing pneumonia associated with pulmonary artery aneurysms in 1285

prevalences of hepatitis A, B, C and E viruses in, 1279 procoagulant mutations and venous thrombosis in, (L) 1298

BELCH J see Ho M

BELEC L see Claudepierre P

BELL AL see Smyth AE BELT EA see Lehtinen JT

BENAMOU P see Seror P

BENICHOU O see Clerc D

BENJAMIN S see Hopkinson N: Hunt IM

BENNETT A. Overview of nimesulide, (S1) 1

BENSON EM see Park KEJ

BERGE E see Clerc D

BERGH K see Mollnes TE

BERGSTRÖM U see Turesson C

BERMAN BM, SINGH BB, LAO L, LANGENBERG P, LI H, HADHAZY V, BARETA J, Hochberg M. A randomized controlled trial of acupuncture as an adjunctive therapy in osteoarthritis of the knee, 346

BESBAS N see Yetgin S

BEUKELMAN CJ see Lafeber FPJG

BHAKTA BB, BRENNAN P, JAMES TE, CHAMBERLAIN MA, NOBLE BA, SILMAN AJ. Behçet's disease: evaluation of a new instrument to measure clinical activity, 728

BHAKTA BB see also Pease CT

BIAGINI P see Gerli R

BIERINGS M. The role of T-cell depletion of autografts for autoimmune diseases, 755

Biggs JC see Snowden JA

BIGONI M see Gallerani M

BIJLSMA JWJ see Blaauw AAM; Jacobs JWG; Lafeber FPJG

BILGIC A see Aksu K

Biological therapies

the emerging role of in the treatment of newly diagnosed rheumatoid arthritis, (S2) 45

in rheumatoid arthritis, health outcomes and health economic issues surrounding the use of (S2) 48

role of in the treatment of refractory rheumatoid arthritis, (S2) 47

Bisson M see Clerc D

BISTONI O see Gerli R

BISWAS A see Handa R
BJARNASON I. Metabolic bone disease in patients with inflammatory bowel disease. (E) 801

BJARNASON I, THJODLEIFSSON B. Gastrointestinal toxicity of non-steroidal anti-inflammatory drugs: the effect of nimesulide compared with naproxen on the human gastrointestinal tract, (S1) 24

BLAAUW AAM, VAN LOON AM, SCHELLEKENS JFP, BIJLSMA JWJ. Clinical evaluation of guidelines and two-test approach for Lyme disease, 1121

BLACK CM see Denton CP

BLANCO JL see Garcia-Pardo G

BLETRY O see Papo T

Blindness in Wegener's granulomatosis, an unusual cause, (L) 578 BLOCKMANS D, MAES A, STROOBANTS S, NUYTS J, BORMANS G, KNOCKAERT

D, BOBBAERS H, MORTELMANS L. New arguments for a vasculitic nature of polymyalgia rheumatica using positron emission tomography, 444 BLYTHE D see McGonagle D

BOBBAERS H see Blockmans D

Body composition changes in pre-menstrual women with systemic lupus erythematosus. Three year follow-up study, 59

Body mass and fat-free body mass, prediction of maximal back muscle strength from indices of, 652

BOERS M. The case for corticosteroids in the treatment of early rheumatoid arthritis, (E) 95

BOERS M see also van der Heijde D; Verhoeven AC

BOLLOW M see Brandt J

BOMBARDIERI S see Swaak AJG

Bone erosion, preliminary study of ultrasound aspiration of in early rheumatoid arthritis, 329

Bone marrow of patients with rheumatoid arthritis, establishment of nurselike stromal cells from: indication of characteristic bone marrow microenvironment, 854

Bone marrow transplantation

allogeneic, transmission of psoriatic arthritis by for chronic myelogenous leukaemia from an HLA-identical donor, (L) 89 autologous, effectiveness and risks of total body irradiation for condi-

tioning in the treatment of autoimmune disease with, 75

Bone mass and bone turnover in polymyalgia rheumatica, (L) 285, 286

Bone, metabolic disease of in patients with inflammatory bowel disease,

Bone mineral density in scleroderma, 371

Rone turnover

and bone mass in polymyalgia rheumatica, (L) 285, 286

relationship to soluble markers of immune activation in postmenopausal women with rheumatoid arthritis, 841

BONGARTZ EB see Drossaers-Bakker KW

BONNARD MN see Perdriger A BONNETTI F see Martini A

BONNOTTE B, KRAUSE D, FANTON A-L, THERON J, CHAUFFERT B, LORCERIE B. False aneurysm of the internal carotid artery in Behcet's disease: successful combined endovascular treatment with stent and coils, (L)

Bonomo L see Giorgi D

BOONEN A see van der Heijde D BORMANS G see Blockmans D BORST HPE see Slaper-Cortenbach ICM BOSSERHOFF AK see Müller-Ladner U

BOST M see Dumolard A

BOT SDM, CASPERS M, VAN ROYEN BJ, TOUSSAINT HM, KINGMA I. Biochemical analysis of posture in patients with spinal kyphosis due to ankylosing spondylitis: a pilot study, 441

BOTTING J see Botting R

BOTTING R, BOTTING J. Directions on rheumatology: past, present and future. Report of a conference to celebrate the 50th anniversary of the Olive Bird Fund held at Churchill College Cambridge, 24-26 September 1998, 887

Вотто M see Naughton MA

Brandt J, Bollow M, Häberle J, Rudwaleit M, Eggens U, Distler A, Sieper J, Braun J. Studying patients with inflammatory back pain and arthritis of the lower limbs clinically and by magnetic resonance imaging: many, but not all, patients with sacoiliitis have spondyloarthropathy, 831

BRANIMIR A see Swaak AJG BRANTUS JF see Chevrel G

BRAUN J see Brandt J; Wilkinson NZ; Yin Z

BRAZIER JE, HARPER R, MUNRO J, WALTERS SJ, SNAITH ML. Generic and condition-specific outcome measurements for people with osteoarthritis of the knee, 870

Breban M see El Maghraoui A

BREEDVELD FC. Future trends in the treatment of rheumatoid arthritis: cytokine targets, (S2) 11

BREEDVELD FC see also Brooks P; Huizinga TWJ; Kraan MC; Tak PP; van den Borne BEE; van der Horst-Bruinsma

BRENNAN FM. Interleukin 10 and arthritis, (E) 293

BRENNAN P see Bhakta BB

BRESNIHAN B see Cunnane G; Shanahan EM

BRIGANTI EM see Kipen Y

BRINKMAN DMC see Vossen JM

BROOKS P, EMERY P, EVANS JF, FENNER H, HAWKLEY CJ, PATRONO C, SMOLEN J, BREEDVELD F, DAY R, DOUGADOS M, EHRICH EW, GUON-BAÑOS J, KVIEN TK, VAN RUSWIJK MH, WARNER T, ZEIDLER H. Interpreting the clinical significance of the differential inhibition of cyclooxygenase-1 and cyclooxygenase-2, 779

BROOKS PM see Snowden JA

BROWNLEE M see Pritchard MH

BRUCE IN see Smyth AE

BRUCKDORFER KR see Denton CP BRUCKNER FE see Johnson SM

BRÜHLMANN P see Leu AJ; Ruof J

BRUN JG see Mollnes TE

BSR Research and Training Committee see Gordon C

BUDD K see Dessein PH

BUETTNER R see Müller-Ladner U BUNCE TD see Denton CP

BUONCOMPAGNI A see Ruperto N
BURMESTER GR, WILSKE KR. Workshop summary: the role of biologicals in the treatment of patients with refractory rheumatoid arthritis, (S2) 47 BURMESTER GR see also Riemekasten G

Bursitis, and destructive joint lesions in idiopathic hypereosinophilic syndrome, (L) 185

BURT RK see Traynor A
BUTLER R see McGeoch P

BÜYÜKBABANI N see Gül

Cox N see Solymossy C CABEZAS JA see Villaverde V CACOUB P see Généreau T CAILLAT-VIGNERON N see Généreau T CAILLEUX N see Marie I CALIN A see Dougados M

CALIN A, NAKACHE J-P, GUEGUEN A, ZEIDLER H, MIELANTS H, DOUGADOS M. Defining disease activity in ankylosing spondylitis: is a com of variables (Bath Ankylosing Spondylitis Disease Activity Index) an appropriate instrument?, 878

CAPELL HA see Morrison E

CARBONE L, TYLAVSKY F, WAN J, MCKOWN K, CHENG S. Bone mineral density in scleroderma, 371

CARR AJ see Dechow E

CARTER N see Williamson L

Cartilage, human articular

in joints of the lower limb. The 'instantaneous' compressive modulus of, 124

ultrasonic measurement of the thickness of in situ, 1269

CASADEMONT J see Miró O

CASEY EB see Flanagan N

CASPERS M see Bot SDM

CASTAGNETO C see Filaci G CATENACCIO M see Cavallo G

CATS HA see Posthumus MD

CAVALLO G, SABADINI L, ROLLO L, CATENACCIO M, LORENZINI S, PIPITONE N, MARCOLONGO R. Nitric oxide synthesis in peripheral blood mononuclear and polymorphonuclear cells from patients with systemic sclerosis. (L) 1301

CD5 and CD23 expression on B cells in peripheral blood and synovial fluid of rheumatoid arthritis patients: relationship with interleukin-4, soluble CD23 and tumour necrosis factor alpha levels, 325

CD20, remission of inflammatory arthropathy in association with anti-CD20 therapy for non-Hodgkin's lymphoma, 1150

CD30, soluble, in early rheumatoid arthritis as a predictor of good response to second-line therapy, 1282

CEDOZ JP see Toussirot E

ČEPONIS A, HIETANEN J, TAMULAITIENÈ M, PARTSCH G, PÄTIÄLÄ H, KONTTINEN YT. A comparative quantitative morphometric study of cell apoptosis in synovial membranes in psoriatic, reactive and rheumatoid arthritis, 431

CHALEM P. GHNASSIA A-M. NORDMANN Y. MENKES CJ. Porphyria cutanca tarda affecting a rheumatoid arthritis patient treated with methotrexate: association or coincidence?, 453

CHALES G see Perdriger A

CHAMBERLAIN MA see Bhakta BB

CHAMOT AM see So A CHAN TM see Kung AWC

CHANDLER C see Walker DJ

Changing times, (E) 1

CHAOUI R see Dörner T

CHARD J see Dieppe P

CHATTERS SJ see O'Duffy EK CHATTOPADHYAY C see Roy DK

CHAUFFERT B see Bonnotte B

CHAUSSADE S see El Maghraoui A

CHECA A see Garcia-Nieto R

CHEN JR, TAKAHASHI M, SUZUKI M, KUSHIDA K, MIYAMOTO S, INOUE T. Comparison of the concentrations of pentosidine in the synovial fluid, serum and urine of patients with rheumatoid arthritis and osteoarthritis, 1275

CHEN WP see Yang LY

CHENEVAL D see Dawson J

CHENG S see Carbone L

CHEONG SK see Fadilah SAW

CHÉRIN P see Généreau T

CHERNAJOVSKY Y. Gene transfer as therapy for rheumatoid arthritis: why, what and how? (E) 804 CHEUNG NT see Mattey DL

CHEVALIER X see Claudepierre P

CHEVREL G, BRANTUS JF, SAINTE-LAUDY J, MIOSSEC P. Allergic pancytopenia to trimethoprim-sulphamethoxazole for Pneumocystis carinii pne following methotrexate treatment for rheumatoid arthritis, (L) 475

Childhood Behçet's disease: clinical features and comparison with adultonset disease, 457

musculoskeletal conditions in, attending two Togolese hospitals, 1010 rationale for high-dose cyclophosphamide and medium-dose total body irradiation in the conditioning of, with progressive systemic and polyarticular juvenile chronic arthritis before autologous stem cell transplantation, 762

with rheumatic diseases, prognosis in: justification for consideration of new therapies, 739

CHIPPS DR see Park KEJ

CHIZZOLINI C see Wicki J

Chlamydia pneumoniae as a triggering infection in reactive arthritis, 411

Chromosomal analysis of peripheral lymphocytes of patients before and after radiation synovectomy with samarium-153 particulate hydroxyapatite, 316

Chronic widespread pain, prevalence and associated features of in the community using the 'Manchester' definition, 275

CHRZANOWSKA-LIGHTOWLERS Z see Szczepanska K CHWALINSKA-SADOWSKA H see Swaak AJG

CIKES N see Swaak AJG

CIOVANNI STRANO C see Ruperto N

CLARKE AE see Sutcliffe N

CLAUDEPIERRE P. ALLANORE Y, BELEC L, LARGET-PIET B, ZARDI L, CHEVALIER X. Increased Ed-B fibronectin plasma levels in spondyloarthropathies: comparison with rheumatoid arthritis patients and a healthy population, 1099

CLERC D, BERGE E, BENICHOU O, PAULE B, QUILLARD J, BISSON M. An unusual case of pigmented villonodular synovitis of the spine: benign,

aggressive and/or malignant?, (L) 476

CLUNIE GPR, WILKINSON ID, LUI D, HALL-CRAGGS MA, PALEY MN, EDWARDS JCW, ELL PJ. Changes in articular synovial lining volume measured by magnetic resonance in a randomized, double-blind, controlled trial of intra-articular samarium-153 particulate hydroxyapatite for chronic knee synovitis, 113

COAKLEY G, MANOLIOS N, LOUGHRAN JT TP, PANAYI GS, LANCHBURY JS. A Fas promoter polymorphism at position -670 in the enhancer region does not confer susceptibility to Felty's and large granular lymphocyte syndromes, 883

COGAN E see Steinfeld SD

COHEN P see Généreau T

Collagenase, cathepsin B and cathepsin L gene expression in the synovial membrane of patients with early inflammatory arthritis, 34

COLLINS DA see Johnson SM

Combination therapy in rheumatoid arthritis (L) 577, 789

Complementary medicine in rheumatic diseases, research on: the need for better quality studies and reproduction of claimed positive results, (E) 387

CONDE A see Fernández-Gutiérrez B

COOPER RG see Fraser A; Mannion AF

COPE AP see Hall FC

Coronary dissection associated with hepatitis C virus-related cryoglobutinaemia, (L) 1299

CORRIGALL VM, GARYFALLOS A, PANAYI GS. The relative proportions of secreted interleukin-2 and interleukin-10 determine the magnitude of rheumatoid arthritis T-cell proliferation to the recall antigen turberculin purified protein derivative, 1203

CORTÉS I see Villaverde V

Corticosteroid(s)

-induced osteoporosis, use of prophylaxis for in hospital practice, (L) 577

the case for in the treatment of early rheumatoid arthritis, (E) 95 future strategies for therapy, 907

local injection, frequency of sepsis after (an inquiry on 1 160 000 injections in rheumatological private practice in France), 1272 in rheumatoid arthritis-the case against, (E) 97

CORTIS E see Ruperto N

Cortisol production, is there a defect in in rheumatoid arthritis?, (E) 298 COUGHLAN RJ see Srinivasan U

COURTOIS H see Marie I

CRAWFORD F, ATKINS D, YOUNG P, EDWARDS J. Steroid injection for heel pain: evidence of short-term effectiveness, 974

CRAWFORD F see also Atkins D CREED FH see Ash G

CROFT P see Dieppe P; Ferry S

Cryptosporidium parvum, sacroiliitis associated with in an HLA-B27-negative patient, (L) 288

CUENDE E, GUINEA J, ORTIZ DE URBINA F, TORRABADELLA S, TEMPLADO JA, VESGA JC. Seronegative oligoarthritis in the course of refractory anaemia with excess blasts, (L) 373 Cunnane G, FitzGerald O, Hummel KM, Gay RE, Gay S, Bresnihan

B. Collagenase, cathepsin B and cathepsin L gene expression in the synovial membrane of patients with early inflammatory arthritis, 34 CUTOLO M see Filaci G; Seriolo B; Walker JG

Cyclooxygenase-1 and cyclooxygenase-2, interpreting the clinical significance of the differential inhibition of, 779

Cyclooxygenase-2, human, molecular model of the interaction with nimesulide. (S1) 14

Cyclophosphamide, high-dose, and medium-dose total body irradiation, rationale for in the conditioning of children with progressive systemic and polyarticular juvenile chronic arthritis before autologous stem cell transplantation, 762

Cyclosporin A

and iloprost treatment of systemic sclerosis: clinical results and interleukin-6 serum changes after 12 months of therapy, 992

therapy in rheumatoid arthritis: only strict application of the guidelines for safe use can prevent irreversible renal function loss, 254

production by synovial T cells in rheumatoid arthritis, 202 role of in the pathogenesis of rheumatoid arthritis, (S2) 3

targets. Future trends in the treatment of rheumatoid arthritis, (S2) 11 Th1-type mRNA in rheumatoid arthritis mononuclear cells induced by streptococcal pyrogenic exotoxin A, (L) 1022

DAGG K see Hamilton J

DAIKELER T, GÜNAYDIN I, EINSELE H, KANZ L, KTTER I. Transmission of psoriatic arthritis by allogeneic bone marrow transplantation for chronic myelogenous leukaemia from an HLA-identical donor, (L) 89

DANIS V see Huang QR

DANKERT T see van der Heijde D DANNECKER GE see Niethammer D

DASGUPTA B see Li C

DAVID J, TOWNSEND S, SATHANATHAN R, KRISS S, DORÉ CJ. The effect of acupuncture on patients with rheumatoid arthritis: a randomized, placebo-controlled cross-over study, 864

Davidson BKS, Kelly CA, Griffiths ID. Primary Sjögren's syndrome in

the North East of England: a long-term follow-up study, 245

DAVIES P see Solymossy C

DAVIES RK see Dechow E

Dawes PT see Layton MA; Mattey DL

DAWSON J, ENGELHARDT P, KASTELIC T, CHENEVAL D, MACKENZIE A, RAMAGE P. Effects of soluble interleukin-1 type II receptor on rabbit antigen-induced arthritis: clinical, biochemical and histological assessment, 401

DAY R see Brooks P

DAYER J-M see Wicki J

D'CRUZ D see Smith PR DE BENEDETTI F see Martini A

DE SONNAVILLE PBJ see Jacobs JWG

De VRIES RRP see van der Horst-Bruinsma

De VRIES-VAN ROSSEN A see Slaper-Cortenbach ICM

DECHOW E, DAVIES RK, CARR AJ, THOMPSON PW. A randomized, double-

blind, placebo-controlled trial of sclerosing injections in patients with chronic low back pain, 1255 DEGNAN BA see Szczepanska K

Dehydroepiandrosterone replacement therapy, should it be provided with glucocorticoids?, 488

DEMOLOMBE-RAGUÉ S see Duhaut P DEMOLS P see Steinfeld SD

DENEYS-LAPORTE V see Huizinga TWJ

DENTON CP, BUNCE TD, DORADO MB, ROBERTS Z, WILSON H, HOWELL K, BRUCKDORFER KR, BLACK CM. Probucol improves symptoms and reduces lipoprotein oxidation susceptibility in patients with Raynaud's phenomenon, 309

DEQUEKER J see Armstrong R

DERCHI L see Filaci G

Dermatomyositis

alfuzosin is not associated with, (L) 579

safety of a home exercise programme: a pilot study, 608

DESSEIN PH, SHIPTON EA, BUDD K. Oral low-dose glucocorticosteroids as compared with intravenous methylprednisolone pulses in the treatment of rheumatoid arthritis, (L) 1304

DEVEY ME see Lewison G

DEVLIN J, WAGSTAFF K, ARTHUR V, EMERY P. Granisetron (Kytril) suppresses methotrexate-induced nausea and vomiting among patients with arthritis and is superior to prochlorperazine inflammmatory (Stemetil), 280

DEVLIN J see also Pease CT

Dexfenfluramine, case of systemic sclerosis that developed under use of, (L) 379

DI BATTISTA JA see Pelletier J-P

Diabetes insipidus, Wegener's granulomatosis and, (L) 684

DIAMOND AG see Szczepanska K

DICKENS CM see Ash G

Diclofenac and nimesulide, comparative efficacy and safety in patients with acute shoulder. (\$1) 33

DIEPPE P, BASLER H-D, CHARD J, CROFT P, DIXON J, HURLEY M, LOHMANDER S, RASPE H. Knee replacement surgery for osteoarthritis: effectiveness, practice variations, indications and possible determinants of utilization, 73

DIEPPE PA see Wotton SF

Diet therapy for the patient with rheumatoid arthritis?, (E) 1039

Dukmans BAC see Drossaers-Bakker KW; van den Borne BEE; van der

Directions on rheumatology: past, present and future. Report of a conference to celebrate the 50th anniversary of the Olive Bird Fund held at Churchill College Cambridge, 24-26 September 1998, 887 DIRESKENELI H see Hamurvudan V Disease activity

assessment in rheumatoid arthritis using magnetic resonance imaging: quantification of pannus volume in the hands, (L) 680

defining in ankylosing spondylitis: is a combination of variables (Bath Ankylosing Spondylitis Disease Activity Index) an appropriate instrument?, 878

form, evaluation of the Turkish translation of for Behçet's syndrome,

high at baseline does not prevent a remission in patients with systemic lupus erythematosus, 724

in patients with systemic lupus erythematosus complicated by haemor-rhagic cystitis, does parenteral oestrogen therapy flare up?, (L) 372 relationship with urinary pyridinium cross-links and disease subsets in ankylosing spondylitis, 21

of rheumatoid arthritis, influence of a partially HLA-matched blood transfusion on 53

score, patient-derived can substitute for a physician-derived in clinical research, 48

serum YKL-40 concentrations in patients with rheumatoid arthritis: relation to, 618

DISTLER A see Brandt J
DIXEY J see Middleton J; Solymossy C

DIXON J see Dieppe P Dixon N see Sinha A

Doctors' handwriting-does size matter?, (L) 683

DODD SM see Smith PR

Doğanavşargil E see Aksu K

DOLAN AL, PANAYI GS. Bone turnover and bone mass in polymyalgia rheumatica, (L) 285

DOLAN P see Mannion AF
DOMLIAN Z see Swaak AJG
DONN RP, FARHAN AJ, BARRETT JH, THOMSON W, WORTHINGTON J, OLLIER WER. Absence of association between interleukin 1 alpha and oligoarticular juvenile chronic arthritis in UK patients, 171

DONNELLY S see Huskisson EC DORADO MB see Denton CP

DORÉ CI see David I

DÖRNER T, FEIST E, CHAOUI R, HIEPE F. Enhanced frequency of autoimmune congenital heart block in female offspring, (L) 380

Dothiepin, effects of on subjects with rheumatoid arthritis and depression, 959

DOUGADOS M, GUEGUEN A, NAKACHE J-P, VELICITAT P, VEYS EM, ZEIDLER H, CALIN A. Ankylosing spondylitis: what is the optimum duration of a clinical study? A one year versus a six weeks non-steroidal antiinflammatory drug trial, 235

Dougados M see also Brooks P; Calin A; El Maghraoui A

Dratwianka B see Swaak AJG DREHER K see Müller-Ladner U

Drewes AM. Pain and sleep disturbances with special reference to fibromyalgia and rheumatoid arthritis, (E) 1035

DROSSAERS-BAKKER KW, HAMBURGER HL, BONGARTZ EB, DIKJMANS BAC, VAN SOESBERGEN RM. Sleep apnoea caused by rheumatoid arthritis. (L) 680

DUANCE VC see Wotton SF

DUFF GW see Wilson AG

DUHAUT P, PINÈDE L, DEMOLOMBE-RAGUÉ S, DUMONTET C, NINET J, SEYDOUX D, LOIRE R, PASQUIER J. Giant cell arteritis and polymyalgia rheumatica: are pregnancies a protective factor? A prospective multicentre case-control study, 118

DUMOLARD A, GAUDIN Ph, JUVIN R, BOST M, PEOC'H M, PHELIP X. SAPHO syndrome or psoriatic arthritis? A familial case study, 463

DUMONTET C see Duhaut P DUMOULIN G see Toussirot E

DUVAL C see Marie I

Economic evaluation, does it have anything to offer the rheumatologist?,

Ed-B fibronectin plasma levels, increased in spondyloarthropathies: comparison with rheumatoid arthritis patients and a healthy population, 1099

EDMONDS J see Fransen M; Huang QR EDWARDS J see Atkins D; Crawford F

EDWARDS JCW see Clunie GPR; O'Duffy EK; Pitsillides AA; Protheroe A Efe B see Korkmaz C

EGGENS U see Brandt J; Yin Z

EHRENSTEIN MR. Antinuclear antibodies and lupus: causes and con-sequences, (E) 691

EHRICH EW see Brooks P EINSELE H see Daikeler T

EL MAGHRAOUI A, DOUGADOS M, FRENEAUX E, CHAUSSADE S, AMOR B, Breban M. Concordance between abdominal scintigraphy using techne-tium-99m hexamethylpropylene amine oxime-labelled leucocytes and ileocolonoscopy in patients with spondylarthropathies and without clinical evidence of inflammatory bowel disease, 543

Electromagnetic tracking for kinematics at the ankle joint complex, preliminary study determining feasibility, 1260

ELL PJ see Clunie GPR

EMERY P see Brooks P; Devlin J; McGonagle D; Pease CT; Solymossy C; Wakefield RJ

EMERY P, MARZO H, PROUDMAN S. Management of patients with newly diagnosed rheumatoid arthritis. (S2) 27
EMERY P, PANAYI G, STURROCK R, WILLIAMS B. Targeted therapies in

rheumatoid arthritis: the need for action, (E) 911

ENDERLIN S see Leu AJ

Endocarditis

bacterial, hypertrophic osteoarthropathy associated with, (L) 375 Streptococcus bovis, polytopic spondylodiscitis secondary to, (L) 686

ENGELHARDT P see Dawson J ENGSTRÖM B see Rolf C

Eosinophilic arthritis, 1188

ERENOĞLU E see Korkmaz C

ERIKSEN EF see Glerup H

ERNST E see White AR

ERTAN S, FRESKO I, APAYDIN H, Özekmekçi S, YAZICI H. Extrapyramidal type rigidity in rheumatoid arthritis, 627

Erythema nodosum, cutaneous necrotizing vasculitis and ankylosing spondylitis, (L) 377

ESCHARD JP see Maillefert JF

European Consortium on Rheumatoid Arthritis Families (ECRAF) see Barrera et al

EVANS JF see Brooks P

Evans P see Helliwell PS

Ewing's sarcoma of the ilium mimicking sacroiliitis, (L) 792

FADILAH SAW, CHEONG SK, SHAHDAN S. Extreme hypereosinophilia in a young male patient presenting with palindromic morphoea, (L) 1158

FALCINI F see Ruperto N FANTON A-L see Bonnotte B

FAREWELL V see Sutcliffe N FARHAN AJ see Donn RP

Fas promoter polymorphism at position -670 in the enhancer region does not confer susceptibility to Felty's and large granular lymphocyte

syndromes, 883 Fas/APO-1 gene polymorphism, association of with systemic lupus erythematosus in Japanese, 516

Fas/Apo-1 promoter polymorphism, evaluation of new, in rheumatoid arthritis and systemic lupus erythematosus patients, 645

FASCIOLO D see Seriolo B

FEIST E see Dörner T

FELDMANN M, MAINI RN. The role of cytokines in the pathogenesis of rheumatoid arthritis, (S2) 3

Felty's syndrome, Fas promoter polymorphism at position -670 in the enhancer region does not confer susceptibility to, 883

Female offspring. Enhanced frequency of autoimmune congenital heart block in, (L) 380

Female patients with rheumatoid arthritis. Relationships between serum 17-8-oestradiol and anticardiolipin antibody concentrations in, (L) 1159 FENNER H see Brooks P

FERNANDES J see Pelletier J-P

FERNÁNDEZ-GUTIÉRREZ B, ZAMORANO J, BATLLE E, ALFONSO F, CONDE A, SANCHEZ-HARGUINDEY L. Coronary dissection associated with hepatitis C virus-related cryoglobutinaemia, (L) 1299

FERNÁNDEZ-PRADA M see Villaverde V

FERRY S, PRITCHARD T, KEENAN J, CROFT P, SILMAN A. Estimating the prevalence of delayed median nerve conduction in the general population, (L) (Reply) 688

Fibromyalgia

determinants of WOMAC function, pain and stiffness scores: evidence for the role of low back pain, symptom counts, fatigue and depression in, 355

pain and sleep disturbances with special reference to, (E) 1035 FILACI G, CUTOLO M, SCUDELETTI M, CASTAGNETO C, DERCHI L, GIANROSSI R, ROPOLO F, ZENTLIN P, SULLI A, MURDACA G, GHIO M, INDIVERI F, PUPPO F. Cyclosporin A and iloprost treatment of systemic sclerosis: clinical results and interleukin-6 serum changes after 12 months of therapy, 992

Finasteride: a steroid-unsparing drug?, (L) 285

Finger joint synovitis in rheumatoid arthritis: quantitative assessment by magnetic resonance imaging, 66

FISCHER R see Swaak AJG FITZGERALD DJ see Shah AA

FITZGERALD O see Cunnane G

FLANAGAN N, CASEY EB, WATSON R, BARNES L. Cutaneous polyarteritis nodosa with seronegative arthritis, (L) 1161

FLORESCU A see Johansen JS

FLOWER RJ. Non-steroidal anti-inflammatory drugs: back to the future, (E) 693

FONSECA A see Villaverde V

FORBES J see Hurst NP

FORMIGA F. MOGA I, PAC M, MITJAVILA F, RIVERA A, PUJOL R. High disease activity at baseline does not prevent a remission in patients with systemic lupus erythematosus, 724

FØRRE Ø see Fraser DA; Haugen M

FRANCO AR see Haier J FRANKE S see Oelzner P

Fransen M, Edmonds J. Gait variables: appropriate objective outcome measurements in rheumatoid arthritis, 663

Fransen M, Edmonds J. Reliability and validity of the EuroQol in patients with osteoarthritis of the knee, 807

FRANZECK UK see Leu AJ FRASER A, VALLOW J, PRESTON A, COOPER RG. Predicting normal 'grip' strength for rheumatoid arthritis patients, 521

FRASER DA, THOEN J, RUSTAN AC, FØRRE Ø, KJELDSEN-KRAGH J. Changes in plasma free fatty acid concentrations in rheumatoid arthritis patients during fasting and their effects upon T-lymphocyte proliferation, 948 FRASER DA see also Haugen M

FREEMAN AM, DASGUPTA B. Angio-neurotic oedema associated with methotrexate treatment in rheumatoid arthritis, 908

FREEMONT AJ see Sari-Kouzel H

FRENEAUX E see El Maghraoui A

FRESKO I see Ertan S; Hamuryudan V; Korkmaz C

FRIES JF. How may quality of life for rheumatoid arthritis patients be enhanced by current and future treatments?, (S2) 35

FRYER AA see Layton MA Fu LW see Yang LY

FURST DE. Leflunomide, mycophenolic acid and matrix metalloproteinase inhibitors, (S2) 14

FURST DE, KEYSTONE E, MAINI RN, SMOLEN JS. Recapitulation of the round-table discussion-assessing the role of anti-tumour necrosis factor therapy in the treatment of rheumatoid arthritis, (S2) 50

GAGO F see Garcia-Nieto R
Gait variables: appropriate objective outcome measurements in rheumatoid arthritis, 663

GAL A see Steiner G

GALEZOWSKI N see Généreau T

GALINDO M, KHAMASHTA MA, HUGHES GRV. Splenectomy for refractory thrombocytopenia in the antiphospholipid syndrome, 848

GALLARDO D see Marsal S

GALLERANI M, GOVONI M, MUCINELLI M, BIGONI M, TROTTA F, MANFREDINI R. Seasonal variation in the onset of acute microcrystalline arthritis, 1003

GALLIVAN S see Solymossy C

GARCIA-NIETO R, PÉREZ C, CHECA A, GAGO F. Molecular model of the interaction between nimesulide and human cyclooxygenase-2, (S1) 14 GARCIA-PARDO G, AUGUET JL, BLANCO JL, ARAÚJO O, LORENZO A, RICHART

C. Polytopic spondylodiscitis secondary to Streptococcus bovis endocarditis, (L) 686

GARNERO A see Seriolo B

GARYFALLOS A see Corrigall VM GASTON JSH see Ramage JM

Gastrointestinal toxicity of non-steroidal anti-inflammatory drugs: the effect of nimesulide compared with naproxen on the human gastrointestinal tract, (S1) 24

GAU JM see Miró Ò

GAUDIN P see Maillefert JF

GAUDIN Ph see Dumolard A

GAY RE see Cunnane G

GAY S see Cunnane G: Müller-Ladner U

Gene transfer as therapy for rheumatoid arthritis: why, what and how?, (E) 804

GÉNÉREAU T, LORTHOLARY O, GUILLEVIN L, CACOUB P, GALEZOWSKI N, CHÉRIN P, BABINET P, HERREMAN G, WECHSLER B, COHEN P, HERSON S, CAILLAT-VIGNERON N. Temporal <sup>67</sup>gallium uptake is increased in temporal arteritis, 709

GEORGE A see Roy DK

GEORGE S see Al-Awadhi AM

GERBER H see Weber M

GERLI R, BISTONI O, LUNARDI C, GIACOMELLI R, TOMASSINI C, BIAGINI P, PITZALIS C. Soluble CD30 in early rheumatoid arthritis as a predictor of good response to second-line therapy, 1282

GERLONI V see Ruperto N

GERSTER J-C see So A

Gню M see Filaci G

GHNASSIA A-M see Chalem P

GIACOMELLI R see Gerli R

GIANROSSI R see Filaci G

Giant cell arteritis

are pregnancies a protective factor? A prospective multicentre casecontrol study, 118 epidemiology of biopsy-positive: special reference to cyclic fluctu-

ations, 1208

in a patient with limited cutaneous systemic sclerosis, (L) 479

presenting as lateral medullary syndrome, (L) 188 skeletal muscle mitochondrial function in, 568

GIBBON WW see McGonagle D; Wakefield RJ

GIBBON WW SEE INCOMISED D, WAREHOLD J GILLOT TJ, STRUTHERS GR. Cutaneous necrotizing vasculitis, erythema nodosum and ankylosing spondylitis, (L) 377 GILLOTT TJ, WHALLETT A, ZAPHIROPOULOS G. Oral desensitization in

patients with chronic tophaceous gout and allopurinol hypersensitivity, (L) 85

GIORGI D. BALACCO GABRIELI C. BONOMO L. The association of optic neuropathy with transverse myelitis in systemic lupus erythematosus,

GLEICHER N. Reproductive failure prior to the onset of clinical autoimmune disease, (E) 485

GLERUP H, ERIKSEN EF. Acroparasthesia-a typical finding in vitamin D deficiency, (L) 482

Glucocorticoid receptor activity, effect of nimesulide on in human synovial fibroblasts, (S1) 11

Glucocorticoids

50 years of treatment in rheumatoid arthritis, (E) 100

in the management of rheumatoid arthritis, 6

oral low-dose, as compared with intravenous methylprednisolone pulses in the treatment of rheumatoid arthritis, (L) 1304

should dehydroepiandrosterone replacement therapy be provided with? 488

GODEAU P see Papo T

GOEI HS see van den Borne BEE

GOODACRE JA see Szczepanska K

GOODFELLOW RM see Richards PJ

GORDON C, SALMON M. Postgraduate degrees for rheumatology trainees: an options appraisal of MD, PhD and MSc degrees, 1290

GORDON C see also Salmon M; Sutcliffe N

GORDON M-M, MADHOK R. Recurrent focal myositis, (L) 1295

Gorham-Stout disease (phantom bone) of the shoulder girdle, (L) 904

Gough A see Solymossy C

GOULDING NJ. Future strategies for corticosteroid therapy, 907

GOVONI M see Gallerani M

GRANINGER W see Swaak AJG

Granisetron (Kytril) suppresses methotrexate-induced nausea and vomiting among patients with inflammmatory arthritis and is superior to prochlorperazine (Stemetil), 280 Green M see McGonagle D Grennan DM see Herrick AL; Roy DK

GRETENER SB see Leu AJ

GRIFFITHS ID see Davidson BKS GRMEK M see Tomšič M

GROENIER KH see Winters JC

GROLMS M see Yin Z

Groupe de Recherche sur l'Artérite à Cellules Géantes (GRACG) see Duhaut et al.

GUDBJÖRNSSON B see Torsteinsdóttir I

GUEDJ D see Krause I

GUEGUEN A see Calin A; Dougados M

GUGGNEBUHL P see Perdriger A GUILLEVIN L. Virus-associated vasculitides, (E) 588

GUILLEVIN L see also Généreau T; Wicki J

GUINEA J see Cuende E

GÜL A, ASLANTAS AB, TEKINAY T, KONIÇE M, ÖZÇELİK T. Procoagulant mutations and venous thrombosis in Behçet's disease, (L) 1298

GÜL A, YILMAZBAYHAN D, BÜYÜKBABANI N, LIE JT, TUNACI M, TUNACI A, INANÇ M, Öcal L, ARAL O, KONIÇE M. Organizing pneumonia associated with pulmonary artery aneurysms in Behçet's disease, 1285

GÜMÜŞDIŞ G see Aksu K GÜNAYDIN I see Daikeler T

GUPTA MN see Jackson MS

HÄBERLE J see Brandt J

HADA T see Matsui K

HADDAD MC see Hajj-Ali RA

HADHAZY V see Berman BM

HAGA H-J see Mollnes TE HAIER J, NASRALLA M, FRANCO AR, NICOLSON GL. Detection of mycoplasmal infections in blood of patients with rheumatoid arthritis, 504 HAJEER AH see Mattey DL; Thomas E

HAJJ-ALI RA, UTHMAN IW, SALTI IA, ZAATARI GS, HADDAD MC, NASR

FW. Wegener's granulomatosis and diabetes insipidus, (L) 684

HAKALA M see Myllykangas-Luosujärvi R

HÅKANSSON L see Torsteinsdóttir

HALL FC, COPE AP, PATEL SD, SØNDERSTRUP G. Isolating the molecular suspect: HLA transgenic mice in the study of human autoimmune disease 697

HALL-CRAGGS MA see Clunie GPR HÄLLGREN R see Torsteinsdóttir I

HAMBURGER HL see Drossaers-Bakker KW

HAMILTON J, DAGG K, STURROCK R, ANDERSON J, BANHAM S. Sleep apnoea caused by rheumatoid arthritis, (L) 679

HAMURYUDAN V, FRESKO I, DIRESKENELI H, TENANT MJ, YURDAKUL S, AKOĞLU T, YAZICI H. Evaluation of the Turkish translation of a disease activity form for Behçet's syndrome, 734
HANDA R. AGGARWAL P. BISWAS A, WIG N. WALI JP. Microscopic polyangi-

itis associated with antiphospholipid syndrome, (L) 478

HANLEY SD see Shanahan EM

HANLY AJ see Abinader A

HANNU T, PUOLKKAINEN M, LEIRISALO-REPO M. Chlamydia pneumoniae as a triggering infection in reactive arthritis, 411

HANSEN M see Johansen JS

HANTOUMI IE see Mavragani CP

HAPP T see Niederstadt C

HARBUZ MS, JESSOP DS. Is there a defect in cortisol production in rheumatoid arthritis?, (E) 298

HARDY CJ, PALMER BP, MORTON SJ, MUIR KR, POWELL RJ. Pregnancy outcome and family size in systemic lupus erythematosus: a case-control study, 559

HAREL L see Krause I

HARINGMAN JJ see Kraan MC

HARMS MC, INNES SM, BADER DL. Forces measured during spinal manipulative procedures in two age groups, 267

HARPER R see Brazier JE

HASEGAWA M, SATO S, IHN H, TAKEHARA K. Enhanced production of interleukin-6 (IL-6), oncostatin M and soluble IL-6 receptor by cultured peripheral blood mononuclear cells from patients with systemic sclerosis 612

Наѕнімото Н see Ogasawara Н

HASLER P see Weber M

HASSELL AB see Mattey DL

HASSFELD W see Hueber W

HAUGEN M, FRASER D, FØRRE Ø. Diet therapy for the patient with rheumatoid arthritis?, (E) 1039

HAWKLEY CJ see Brooks P HAY EM see Thomas E

HAYASHI K see Horiuchi T

HAYASHI N see Matsui K HAZES JMW see van der Horst-Bruinsma

Heel pain

steroid injection for: evidence of short-term effectiveness, 974 a systematic review of treatments, 968

HEIM M see Barzilai A

HEIN G see Oelzner P

HEIN R see Müller-Ladner U

HELBERT M see Smith PR

HELIÖVAARA M see Mäkelä M

HELLIWELL PS. HETTHEN J. Primary care rheumatology-leading the way?, (E) 117

HELLIWELL PS, O'HARA M, HOLDSWORTH J, HESSELDEN A, KING T, EVANS P. A (2-month randomized controlled trial of patient education on radiographic changes and quality of life in early rheumatoid arthritis,

HELLIWELL PS see also Hetthen J; Woodburn J

Hepatitis A, B, C and E viruses, prevalence in Behçet's disease, 1279

Hepatitis B vaccination, rheumatic disorders developed after, 978

Hepatitis B virus, successful treatment of polyarteritis nodosa related to, with a combination of lamivudine and interferon alpha, (L) 183

Hepatitis C virus-related cryoglobutinaemia, coronary dissection associated with (L) 1299

HERREMAN G see Généreau T

HERRICK AL, GRENNAN DM, AARONS L. Lack of interaction between methotrexate and penicillins, 284

HERRICK AL see also Sari-Kouzel H

HERSON S see Généreau T

HESLOP PS see Walker DJ

HESSELDEN A see Helliwell PS

HETTHEN J, HELLIWELL PS. A comparison between primary care-led rheumatology services and secondary care provision, (L) 1294

HETTHEN J see also Helliwell PS

HIDALGO V see Villaverde V

HIEPE F see Dörner T; Riemekasten G HIETANEN J see Čeponis A

HIGASHINO K see Matsui K

HIGGENS CS see Schott JM

Нібисні М see Horiuchi Т

HILLS BA see Vecchio P

HIROISHI K see Matsui K HIROSE S see Ogasawara H

HISHIKAWA T see Ogasawara H Histoplasmosis of the wrist: a case report, 906

HIV infection, drug-induced lupus nephritis in, (L) 1017

HIV-infected haemophiliacs versus other risk groups, characteristics of septic arthritis in, 139

HLA associations, clinical, radiographic and, as markers for different patterns of psoriatic arthritis, 332

HLA class I antigen. Molecular mimicry between human endogenous retrovirus clone 4-1 and, with reference to the pathogenesis of systemic lupus erythematosus, (L) 1163

HLA DMB1\*0101-0101 genotype. Positive association with rheumatoid arthritis, 448

HLA, partially matched blood transfusion, influence of on the disease activity of rheumatoid arthritis, 53

HLA transgenic mice in the study of human autoimmune disease. Isolating the molecular suspect, 697

HLA-B27-negative patient. Sacroiliitis associated with Cryptosporidium arvum in an. (L) 288

HLA-DRB1 alleles encoding the DERAA amino acid motif. The influence of on radiological outcome in rheumatoid arthritis, 1221

HLA-identical donor. Transmission of psoriatic arthritis by allogeneic bone marrow transplantation for chronic myelogenous leukaemia from an, (1)89

Ho M, Belch J. Causes of mortality in systemic sclerosis, (L) 283

HOCHBERG M see Berman BM

HODGKINS P, HULL R. Use of prophylaxis for corticosteroid-induced osteoporosis in hospital practice, (L) 577

HOFBAUER LC, KLASSEN RA, KHOSLA S. Gorham-Stout disease (phantom bone) of the shoulder girdle, (L) 904

HOFFMANN U see Leu AJ

Hot. C see Houtman PM

HOLDSWORTH J see Helliwell PS

Home exercise programme in patients with polymyositis and dermatomyositis, safety of: a pilot study, 608

HOOGERBRUGGE PM see Slaper-Cortenbach ICM; Vossen JM

HOPKINSON N, PARHAM D, BENJAMIN S. Silastic prostheses-a forgotten cause

of lymphadenopathy in rheumatoid arthritis, (L) 480 Hoque SN, Minassian M, Clipstone S, Lloyd-Owen SJ, Sheridan E, LESSING MPA. Melioidosis presenting as septic arthritis in Bengali men in East London, (L) 1029

HORDON LD, le GALLEZ P, ISDALE AH. Oral methotrexate: hazard of different tablet strength, (L) 1304

HORIUCHI T, NSHIZAKA H, YASUNAGA S, HIGUCHI M, TSUKAMOTO H, HAYASHI K, NAGASAWA K. Association of Fas/APO-1 gene polymorphism with systemic lupus erythematosus in Japanese, 516

Hormone replacement therapy. Osteopenia in young hypogonadal women with systemic lupus erythematosus receiving chronic steroid therapy: a randomized controlled trial comparing calcitriol and, 1239

HØRSLEV-PETERSEN K see Johansen JS

HOUSSIEN DA, STUCKI G, SCOTT DL. A patient-derived disease activity score can substitute for a physician-derived disease activity score in clinical research 48

HOUTMAN PM, MARCK KW, HOL C. Histoplasmosis of the wrist: a case report, 906

HOWELL K see Denton CP

HUANG QR, DANIS V, LASSERE M, EDMONDS J, MANOLIOS N. Evaluation of a new Apo-1/Fas promoter polymorphism in rheumatoid arthritis and systemic lupus erythematosus patients, 645

HUANG WJS see Yang LY

HUEBER W, HASSFELD W, SMOLEN JS, STEINER G. Sensitivity and specificity of anti-Sa autoantibodies for rheumatoid arthritis, 155

HUGHES AE see Smyth AE

HUGHES GRV, KHAMASHTA MA. Prolactin, the immune response and lupus, (E) 296

HUGHES GRV see also Atsumi T; Galindo M

HUIZINGA TWJ, VAN DER LINDEN MW, DENEYS-LAPORTE V, BREEDVELD FC. Interleukin-10 as an explanation for pregnancy-induced flare in systemic lupus erythematosus and remission in rheumatoid arthritis. 496

HUIZINGA TWJ see also van der Horst-Bruinsma; Verweij CL

HULSMANS HMJ see Jacobs JWG HUMMEL KM see Cunnane G

HUNT IM, SILMAN AJ, BENJAMIN S, MCBETH J, MACFARLANE GJ. The prevalence and associated features of chronic widespread pain in the community using the 'Manchester' definition of chronic widespread pain, 275

HUNT IM see also Niobyu P

HUNT S see Lloyd ME

HURLEY M see Dieppe P

HURST NP. Combination therapy in rheumatoid arthritis, (L) 789

HURST NP, FORBES J. Does economic evaluation have anything to offer the rheumatologist?, (E) 2

HUSKISSON EC, DONNELLY S. Hyaluronic acid in the treatment of osteoarthritis of the knee, 602

HUTCHINSON M see Shanahan EM

Hyaluronic acid in the treatment of osteoarthritis of the knee, 602

Hypereosinophilia, extreme, in a young male patient presenting with palindromic morphoea, (L) 1158

Hypereosinophilic syndrome, idiopathic, destructive joint lesions and bursitis in. (L) 185

Hypertrophic osteoarthropathy associated with bacterial endocarditis, (L)

ICHIKAWA K see Atsumi T

IHN H see Hasegawa M

Iloprost and cyclosporin A treatment of systemic sclerosis: clinical results and interleukin-6 serum changes after 12 months of therapy, 992

Imaging of pelvic fracture in a patient with psoriatic arthritis, (L) 902 Immune activation, soluble markers of, relationship to bone turnover in post-menopausal women with rheumatoid arthritis, 841

Immunohistological analysis of synovial tissue for differential diagnosis in early arthritis, 1074

Immunosuppressive therapy

long-term toxicity of in juvenile rheumatic diseases, 743

normotensive renal failure in a patient with systemic sclerosis and pantineutrophil cytoplasmic autoantibodies which developed into Paget's disease of bone after, (L) 190

side-effects of long-term, versus morbidity in autologous stem cell rescue: striking the balance, 747

IMPIVAARA O see Mäkelä M

INANÇ M see Gül A; Radway-Bright EL

INDIVERI F see Filaci G

INNES SM see Harms MC

INOKUMA S, SETOGUCHI K, OHTA T, MATSUZAKI Y, YOSHIDA A. Serum leucine aminopeptidase as an activity indicator in systemic lupus erythematosus: a study of 46 consecutive cases, 705

INOUE K, SHICHIKAWA K, OTA H. Prevalence of hip osteoarthritis and acetabular dysplasia in Kamitonda: from a longitudinal populationbased epidemiological study of rheumatic diseases in Japan, 793

INOUE T see Chen JR; Naito K

Interferon alpha, successful treatment of polyarteritis nodosa related to hepatitis B virus with a combination of lamivudine and, (L) 183

Interferon beta treatment, effects on arthritis, 362

Interferon gamma, tumour necrosis factor alpha and granulocyte-macrophage colony stimulating factor levels, evaluation of in juvenile chronic arthritis, 468

Interferon gamma- and interleukin-4-producing T cells, quantification of in synovial fluid and peripheral blood of arthritis patients, 214

Interferon gamma-/interleukin-4-positive T cells, elevated levels of found in synovial fluid and synovial membrane of rheumatoid arthritis patients can be changed by interleukin-4 but not by interleukin-10 or transforming growth factor beta, 1058

Interferon-4, soluble CD23 and tumour necrosis factor alpha levels Relationship with CD5 and CD23 expression on B cells in peripheral blood and synovial fluid of rheumatoid arthritis patients, 325

Interleukin-1 alpha, absence of association between, and oligoarticular juvenile chronic arthritis in UK patients, 171

Interleukin-1 type II receptor, soluble, effects of on rabbit antigen-induced arthritis: clinical, biochemical and histological assessment, 401

Interleukin-2 and interleukin-10, relative proportions of secreted, determine the magnitude of rheumatoid arthritis T-cell proliferation to the recall antigen turberculin purified protein derivative, 1203

Interleukin-4-/interferon gamma-positive T cells, elevated ratio of found in synovial fluid and synovial membrane of rheumatoid arthritis patients can be changed by interleukin-4 but not by interleukin-10 or transforming growth factor beta, 1058

Interleukin-4-producing T cells in synovial fluid and peripheral blood of arthritis patients. Quantification of interferon gamma-and, 214 Interleukin-6

oncostatin M and soluble IL-6 receptor, enhanced production of by cultured peripheral blood mononuclear cells from patients with systemic sclerosis, 612

serum changes, cyclosporin A and iloprost treatment of systemic sclerosis: clinical results and, after 12 months of therapy, 992

Interleukin-10

as an explanation for pregnancy-induced flare in systemic lupus erythematosus and remission in rheumatoid arthritis, 496 and arthritis, (E) 293

Interleukin-10 receptor cDNA gene (IL10R), polymorphisms within, in Japanese patients with systemic lupus erythematosus, 1142

IOANNIDIS JPA see Mavragani CP

IOANNOU Y see Sultan SM

ISDALE AH. Sacral insufficiency fractures: an unsuspected cause of low back pain, (L) 90

ISDALE AH see also Hordon L.D.

ISENBERG D, RAMSEY-GOLDMAN R. Assessing patients with lupus: towards a drug responder index, 1045

ISENBERG D see also Ames PRJ; Radway-Bright EL; Sultan SM; Sutcliffe N Italian Pediatric Rheumatology Study Group see Ruperto et al IWAHASHI T see Muro H

JACKI SH, UHL M, ADLER CP, PETER HH, von KEMPIS J. Predominant ankle arthropathy in hereditary haemochromatosis, (L) 378 JACKSON C see Middleton J

JACKSON MS, BAGG J, GUPTA MN, STURROCK RD. Oral carriage of staphylococci in patients with rheumatoid arthritis, 572

JACOBS JWG, DE SONNAVILLE PBJ, HULMANS HMJ, VAN RINSUM AC,

BIJLSMA JWJ. Polyarticular heterotopic ossification complicating critical illness, 1145

JACOBS LD see Rolf C

JACOBSSON L see Turesson C

JAMES D see Solymossy C

JAMES TE see Bhakta BB

JAMES WH. The rates of alcohol-related deaths from accidents and violence in subjects with ankylosing spondylitis and rheumatoid arthritis, (L) 91  $_{\mbox{\scriptsize JANSEN}}$  TLThA see Laan RFJM

JÄNTTI J, AHO K, KAARELA K, KAUTIAINEN H. Work disability in an cohort of patients with seropositive rheumatoid arthritis: a 20 year study, 1138 JARRETA D see Miró Ò

JASSIM A, KUMAR N, KELLY C. Adult Still's disease with nephrotic syndrome at presentation, (L) 283

JAYSON MI see Ash G; Sari-Kouzel H

JESSOP DS see Harbuz MS

JIMENEZ-MAZUECOS J, YEBRA-BANGO M, SANCHEZ-RUIS A, VILLARREAL-GARCIA-LOMAS M. Weber-Christian disease associated with lupus anticoagulant and anticardiolipin antibodies, (L) 796

JOHANSEN JS, STOLTENBERG M, HANSEN M, FLORESCU A, HØRSLEV-PETERSEN K, LORENZEN I, PRICE PA. Serum YKL-40 concentrations in patients with rheumatoid arthritis: relation to disease activity, 618

JOHANSSON H see Petursdottir V

JOHNSON SM, PATEL S, BRUCKNER FE, COLLINS DA. 5'-Nucleotidase as a marker of both general and local inflammation in rheumatoid arthritis patients, 391

destruction, involvement of fibroblast growth factor-2 in of rheumatoid arthritis patients, 714

destructive lesions and bursitis in idiopathic hypereosinophilic syndrome, (L) 185

detection of DNA from a range of bacterial species in, of patients with a variety of arthritides using a nested, broad-range polymerase reaction chain, 260

detection of mycobacteria in samples from patients with arthritis using a genus-specific polymerase chain reaction and sequence analysis,

immobilization reduces synovial fluid hyaluronan concentration and is accompanied by changes in the synovial intimal cell populations,

of the lower limb, 'instantaneous' compressive modulus of human articular cartilage in, 124

JONES AKP. Gastrointestinal tolerability of meloxicam compared to diclofenac in osteoarthritis patients, (L) 793

JONES HW see Wilkinson NZ

JONES PW see Layton MA; Mattey DL

JONKER M see Tak PP JÓNSSON H see Qvindesland A

JORDANOGLOU J see Tatsis G

JOUNIEAUX V see Nevez G

Juvenile chronic arthritis

evaluation of tumour necrosis factor alpha, interferon gamma and granulocyte-macrophage colony stimulating factor levels in, 468

oligoarticular, absence of association between interleukin 1 alpha and, in UK patients, 171

refractory, autologous stem cell transplantation 737, 764

proposed guidelines, 777

responsiveness of outcome measurements in, 176

Juvenile rheumatic diseases, long-term toxicity of immune suppression in, 743 JUVIN R see Dumolard A; Maillefert JF

KAARELA K see Jäntti J; Lehtinen JT KABASAKAL Y see Aksu K KABUKÇUOĞLU S see Korkmaz C KALDEN JR see Paulus HE KALDERN JR see Swaak AJG

KANEKO H see Ogasawara H

KANEKO M see Tomita T

KANEKO Y see Ogasawara H

KANZ L see Daikeler T

KARAAGAOGLU E see Yetgin S

KASKANI E see Tektonidou M KASTELIC T see Dawson J

KATO T see Muro H

KAUPPI MJ see Lehtinen JT

KAUTIAINEN H see Jäntti J; Lehtinen JT; Myllykangas-Luosujärvi R

KAVANAUGH A see Lipsky PE KAWAGUCHI H see Manabe N

Kaznowska Z see Smoleńska Z Keat A see Schott JM

KEENAN J see Ferry S

KEINO H see Maeda T

KELLY C see Davidson BKS; Jassim A

KESER G see Aksu K

Ketoprofen, intra-articular absorption and distribution of after topical plaster application and oral intake in 100 patients undergoing knee arthroscopy, 564

KEYSTONE E see Furst DE

KHAMASHTA MA see Atsumi T; Galindo M

KHOSLA S see Hofbauer LC

KIDD E see Walker DJ

KING T see Helliwell PS

KINGMA I see Bot SDM KINGSLEY GH see Wilkinson NZ

KIPEN Y, BRIGANTI EM, STRAUSS BJG, LITTLEJOHN GO, MORAND EF. Three year follow-up of body composition changes in pre-menstrual women with systemic lupus erythematosus, 59

KIRWAN JR, BÁLINT G, SZEBENYI B. Anniversary: 50 years of glucocorticoid treatment in rheumatoid arthritis, (E) 100

Kiss E see Swaak AJG

KJELDSEN-KRAGH J see Fraser DA

KLARLUND M, Østergaard M, LORENZEN I. Finger joint synovitis in rheumatoid arthritis: quantitative assessment by magnetic resonance imaging, 66

KLASSEN LW see O'Dell JR

KLASSEN RA see Hofbauer LC

Klebsiella serotypes K26, K36 and K50, increased antibody responses in patients with ankylosing spondylitis, (L) 481

Knee arthroscopy, intra-articular absorption and distribution of ketoprofen after topical plaster application and oral intake in 100 patients undergoing, 564

Knee osteoarthritis

acupuncture as an adjunctive therapy. A randomized controlled

femoral intercondylar notch measurements in, 554

generic and condition-specific outcome measurements, 870

measurement of matrix metalloproteinases (MMPs) and tissue inhibitor of metalloproteinases-1 (TIMP-1): comparison with generalized osteoarthritis, 510

reliability and validity of the EuroQol in, 807

replacement surgery for: effectiveness, practice variations, indications and possible determinants of utilization, 73

Knee synovitis, chronic, changes in articular synovial lining volume measured by magnetic resonance in a randomized, double-blind, controlled trial of intra-articular samarium-153 particulate hydroxyapatite, 113

KNEKT P see Mäkelä M

KNOCKAERT D see Blockmans D

KOIKE T see Atsumi T

Koniçe M see Gül A Konttinen YT see Čeponis A

KORKMAZ C, EFE B, TAL N, KABUKÇUOĞLU S, ERENOĞLU E. Sarcoidosis with palpable nodular myositis: periostitis and large-vessel vasculitis simulating Takayasu's arteritis, (L) 287

KORKMAZ C, FRESKO I, YAZICI H. A case of systemic sclerosis that developed under dexfenfluramine use, (L) 379

KORPELA M see Pertovaara M

Kostense P see van der Heijde D

KÖTTER I see Daikeler T

Kouri T see Pertovaara M

Kovacs A see Swaak AJG

Kovacs L see Swaak AJG

Kowal-Bielecka O see Leu AJ

KRAAN MC, HARINGMAN JJ, POST WJ, VERSENDAAL J, BREEDVELD FC, TAK PP. Immunohistological analysis of synovial tissue for differential diagnosis in early arthritis, 1074

KRAAN MC see also Tak PP

KRAUSE D see Bonnotte B

Krause I, Uziel Y, Guedi D, Mukamel M, Harel L, Molad Y, Weinberger A. Childhood Behçet's disease: clinical features and comparison with adult-onset disease, 457

KRISS S see David J

KROOT EJA see van de Putte LBA

KUGA Y see Manabe N

KUIS W, WULFFRAAT NM, PETTY RE. Autologous stem cell transplantation: an alternative for refractory juvenile chronic arthritis, 737

Kuis W see also Wulffraat NM

KUMAR N see Jassim A

KÜMMERLE-DESCHNER J see Niethammer D

KUNAVER M see Steiner G

KUNG AWC, CHAN TM, LAU CS, WONG RWS, YEUNG SSC. Osteopenia in young hypogonadal women with systemic lupus erythematosus receiving chronic steroid therapy: a randomized controlled trial comparing calcitriol and hormone replacement therapy, 1239

KUNTZ JL see Maillefert JF

KUPER HH see Albers JMC

KUSHIDA K see Chen JR; Naito K; Takahashi M

KVEDER T see Tomšič M

KVIEN TK see Brooks P

LA CORTE R see Trotta F

LAAN RFJM, JANSEN TLThA, VAN RIEL PLCM. Glucosteroids in the management of rheumatoid arthritis, 6

LAFEBER FPJG, BEUKELMAN CJ, VAN DEN WORM E, VAN ROY JLAM, VIANEN ME, VAN ROON JAG, VAN DUK H, BULSMA JWJ. Apocynin, a plant-derived, cartilage-saving drug, might be useful in the treatment of rheumatoid arthritis, 1088

LAGAAY EM see van der Horst-Bruinsma

LAMBERT M see Atkins D

LANCHBURY JS see Coakley G

LANDEWÉ RBM see van den Borne BEE

LANG B see Müller-Ladner U

LANGENBERG P see Berman BM

LAO L see Berman BM LARGET-PIET B see Claudepierre P

Larsen scoring of digitalized X-ray images, 1127

LASSERE M see Huang QR

LAU CS see Kung AWC

LAWSON EK see Mijiyawa M

LAXER RM. Long-term toxicity of immune suppression in juvenile rheumatic

LAYTON MA, JONES PW, ALLDERSEA JE, STRANGE RC, FRYER AA, DAWES PT, MATTEY DL. The therapeutic response to D-penicillamine in rheumatoid arthritis: influence of glutathione S-transferase polymorphisms, 43

Le Gallez P see Hordon LD Le Liboux A see Rolf C

LE Thi HUONG D see Papo T

LECOQ D'ANDRE F see Seror P

Leflunomide, mycophenolic acid and matrix metalloproteinase inhibitors, (S2) 14

LEHTINEN JT, LEHTO MUK, KAARELA K, KAUTIAINEN HJ, BELT EA, KAUPPI MJ. Radiographic joint space in rheumatoid acromioclavicular joints: a 15 year prospective follow-up study in 74 patients, 1104

LEHTO MUK see Lehtinen JT

LEIRISALO-REPO M see Hannu T

LEPORE L see Ruperto N

LEPRETRE S see Marie 1

LEU AJ, GRETENER SB, ENDERLIN S, BRÜHLMANN P, MICHEL BA, KOWAL-BIELECKA O, HOFFMANN U, FRANZECK UK. Lymphatic microangiopathy of the skin in systemic sclerosis, 221

LEVESQUE H see Marie I

Lewison G, Devey ME. Bibliometric methods for the evaluation of arthritis research, 13

Li C, Dasgupta B. Sacral insufficiency fractures: reply to correspondence, (L) 903

Li H see Berman BM

LIAROPOULOS L. Economic comparison of nimesulide and diclofenac, and the incidence of adverse events in the treatment of rheumatic disease in Greece, (S1) 39

LIE JT see Gül A

LIENCE E see Marsal S

LIMBURG PC see Posthumus MD

LIN CY see Yang LY

LIOZON E, LY K, AUBARD Y, VIDAL E. Intravenous immunoglobulins for adult Still's disease and pregnancy, (L) 1024

Liposomal clodronate eliminates synovial macrophages, reduces inflammation and ameliorates joint destruction in antigen-induced arthritis, 818 LIPSKY PE, KAVANAUGH A. The impact of pharmaco-economic considera-

tions on the utilization of novel anti-rheumatic therapies, (S2) 41 LITTLEJOHN GO see Kipen Y; Walker JG

LIU L see Yin Z

LLOYD DC see O'Duffy EK

LLOYD ME, HUNT S, SPECTOR TD. Imaging of pelvic fracture in a patient with psoriatic arthritis, (L) 902

LOCATELLI F see Martini A

Logar D see Swaak AJG; Tomšič M

LOHMANDER S see Dieppe P LOIRE R see Duhaut P

LORCERIE B see Bonnotte B; Maillefert JF

LORENZEN I see Johansen JS; Klarlund M

LORENZINI S see Cavallo G

LORENZO A see Garcia-Pardo G LORTHARY O see Généreau T

LOUGHRAN JR TP see Coakley G

Low back pain

chronic. A randomized, double-blind, placebo-controlled trial of sclerosing injections, 1255

sacral insufficiency fractures as an unsuspected cause, (L) 90, 903

LOZA E, TINTURÉ T, SÁNCHEZ-IBARROLA A. CD5 and CD23 expression on B cells in peripheral blood and synovial fluid of rheumatoid arthritis patients: relationship with interleukin-4, soluble CD23 and tumour necrosis factor alpha levels, 325

LOZADA CJ see Abinader A

Lui D see Clunie GPR

LUNARDI C see Gerli R

LUNDBERG I see Alexanderson H

Lung function in Adamantiades- Behçet's disease, (L) 1018

Lupus

antinuclear antibodies and: causes and consequences, (E) 691 assessing patients: towards a drug responder index, 1045 see also Systemic lupus erythematosus

Lupus anticoagulant and anticardiolipin antibodies, Weber-Christian disease associated with, (L) 796

Lupus nephritis, drug-induced, in HIV infection, (L) 1017

Ly K see Liozon E

Lyme disease, clinical evaluation of guidelines and two-test approach for, 1121

Lymphatic microangiopathy of the skin in systemic sclerosis, 221

Lymphocyte syndromes, large granular, Fas promoter polymorphism at position -670 in the enhancer region does not confer susceptibility to. 883

MACCARIO R see Martini A

MACFARLANE G see Hunt IM; Njobvu P

MACGREGOR AJ, SPECTOR TD. Twins and the genetic architecture of osteoarthritis, (E) 583

MACKENZIE A see Dawson J

MACLENNAN K see Protheroe A

Macrophage-colony stimulating factor (M-CSF), high serum level of in Adult Still's disease, (L) 477

Марнок R see Gordon M-M

MAEDA T, KEINO H, ASAHARA H, TANIGUCHI M, NISHIOKA K, SUMIDA T. Decreased TCR AV24AJ18\* double-negative T cells in rheumatoid synovium, (L) 186

MAES A see Blockmans D

Magnetic resonance imaging

assessment of disease activity in rheumatoid arthritis: quantification of pannus volume in the hands, (L) 680

quantitative assessment of finger joint synovitis in rheumatoid arthritis, 66

studying patients with inflammatory back pain and arthritis of the lower limbs clinically and by: many, but not all, patients with sacroillitis have spondyloarthropathy, 831

MAILLEFERT JF, SIBILIA J, TOUSSIROT E, VIGNON E, ESCHARD JP, LORCERIE B, JUVIN R, PARCHIN-GENESTE N, PIROTH C, WENDLING D, KUNTZ JL, TAVERNIER C, GAUDIN P. Rheumatic disorders developed after hepatitis

B vaccination, 978 Maini RN, Wolfe F. Workshop summary: health outcomes and health economic issues surrounding the use of biological therapies in patients with rheumatoid arthritis, (S2) 48

MAINI RN see also Feldmann M; Furst DE

MAKDASSI R see Nevez G

MÄKELÄ M, HELÏVAARA M, SAINIO P, KNEKT P, IMPIVAARA O, AROMAA A. Shoulder joint impairment among Finns aged 30 years or over: prevalence, risk factors and co-morbidity, 656

Manabe, Oda H, Nakamura K, Kuga Y, Uchida S, Kawaguchi H. Involvement of fibroblast growth factor-2 in joint destruction of rheumatoid arthritis patients, 714

MANFREDINI R see Gallerani M

MANGER K see Swaak AJG

MANNION AF, ADAMS MA, COOPER RG, DOLAN P. Prediction of maximal

back muscle strength from indices of body mass and fat-free body mass. 652

Manolios N see Coakley G; Huang QR

MARTENSSON U see Mollnes TE

MARCHESONI A see Swaak AJG

MARCK KW see Houtman PM MARCOLONGO R see Cavallo G

MARCUSON RW see Sari-Kouzel H

MARIE I, LEVESQUE H, CAILLEUX N, LEPRETRE S, DUVAL C, TILLY H, COURTOIS H. An uncommon association: Sjögren's syndrome and autoimmune myelofibrosis, (L) 370

MARSAL S, ARMADANS-GIL L, MARTINEZ M, GALLARDO D, RIBERA A, LIENCE E. Clinical, radiographic and HLA associations as markers for different patterns of psoriatic arthritis, 332

MARTEL-PELLETIER J see Pelletier J-P

MARTIN-MOLA E see Villaverde V

MARTINEZ M see Marsal S

Martini A, Maccario R, Ravelli A, Montagna D, De Benedetti F, Bonetti F, Viola S, Zecca M, Perotti C, Locatelli F. Marked and sustained improvement 2 years after autologous stem cell transplantation in a girl with systemic sclerosis, 773

MARTINI A see also Ruperto N MARTINOWITZ U see Barzilai A

MARUYAMA N see Ogasawara H

MARZO H see Emery P

Matrix metalloproteinase inhibitors, leflunomide, mycophenolic acid and, (S2) 14

Matrix metalloproteinase-3 (MMP-3)

serum levels in relation to the development of radiological damage in patients with early rheumatoid arthritis, 1081

serum, in rheumatoid arthritis: correlation with systemic inflammation but not with erosive status, 407

Matrix metalloproteinases (MMPs)

autocrine induction of gliostatin/platelet-derived endothelial cell growth factor (GLS/PD-ECGF) and GLS-induced expression of in rheumatoid arthritis synoviocytes, 1195

and tissue inhibitor of metalloproteinases-1 (TIMP-1), measurement of in patients with knee osteoarthritis: comparison with generalized osteoarthritis, 510

MATSUI K, TSUCHIDA T, HIROISHI K, TOMINAGA K, HAYASHI N, HADA T, HIGASHINO K. High serum level of macrophage-colony stimulating factor (M-CSF) in adult-onset Still's disease, (L) 477

MATSUI N see Muro H

MATSUZAKI Y see Inokuma S

MATTEY DL, HASSELL AB, PLANT MJ, CHEUNG NT, DAWES PT, JONES PW, THOMSON W, POULTON KV, HAJEER AH, OLLIER WER. The influence of HLA-DRB1 alleles encoding the DERAA amino acid motif on radiological outcome in rheumatoid arthritis, 1221

MATTEY DL see also Layton MA

MAVRAGANI CP, IOANNIDIS JPA, TZIOUFAS AG, HANTOUMI IE, Moursopoulos HM. Recurrent pregnancy loss and autoantibody pro-file in autoimmune diseases, 1228

McBeth J see Hunt IM

McDonald K see Shanahan EM

McGeer EG see Waterfield JD McGeer PL see Waterfield JD

McGeoch P, Butler R. The ARC steroid trial and its effect on clinical practice, (L) 905

McGonagle D, Gibbon W, O'Connor P, Blythe D, Wakefield R, Green M, VEALE D, EMERY P. A preliminary study of ultrasound aspiration of bone erosion in early rheumatoid arthritis, 329

McKown K see Carbone L McMurray NE see Walker JG

McNally JD see Schott JM McNally R, Rooney B, Stewart O. Re: Tench et al. An insight into rheumatology resources available on the World Wide Web, (L) 1027 Median nerve conduction, delayed, estimating prevalence in the general

population, (L) 688

Mediterranean fever, familial, and polyarteritis nodosa complicated by spontaneous bilateral perirenal and subcapsular splenic haemorrhage, (L) 794

Melioidosis presenting as septic arthritis in Bengali men in East London,(L) 1029

MELSOM RD. Familial hypersensitivity to allopurinol with subsequent desensitization, (L) 1301

MENKES CJ see Chalem P

MERT A see Basaranoglu M

Metabolic bone disease in patients with inflammatory bowel disease, (E) 801 Methotrexate

-induced nausea and vomiting, granisetron (Kytril) suppresses, among patients with inflammmatory arthritis and is superior to prochlorperazine (Stemetil), 280

allergic pancytopenia to trimethoprim-sulphamethoxazole for nocystis carinii pneumonia following treatment for rheumatoid arthritis, (L) 475

angio-neurotic oedema associated with treatment in rheumatoid arthritis, 908

effect on blood purine and pyrimidine levels in patients with rheumatoid arthritis, 997

lack of interaction between, and penicillins, 284

oral, hazard of different tablet strengths, (L) 382, 1304

porphyria cutanea tarda affecting a rheumatoid arthritis patient treated with: association or coincidence?, 453

Methylprednisolone, intravenous pulses as compared with oral low-dose glucocorticoids in the treatment of rheumatoid arthritis, (L) 1304 MEYBOOM-DE Jong B see Winters JC

MEYER R see Riemekasten G

MIA (melanoma inhibitory activity): a potential serum marker for rheumatoid arthritis, 148

MICHEL BA see Leu AJ; Ruof J

Microscopic polyangiitis associated with antiphospholipid syndrome, (L) 478 Microvascular abnormalities in Sjögren's syndrome: nailfold capillaroscopy, 826

MIDDLETON J, WHITE S, PARRY E, JACKSON C, DIXEY J, ASHTON B. Changes in serum chondroitin sulphate epitopes 3-B-3 and 7-D-4 in early rheumatoid arthritis, 837

MIELANTS H see Calin A

MIGLIAVACCA D see Ruperto N

MIJIYAWA M, ONIANKITAN O, ATTOH-MENSAH K, TÉKOU A, LAWSON EK, PRIULI GB, ASSIMADI JK. Musculoskeletal conditions in children attending two Togolese hospitals, 1010

MIOSSEC P see Chevrel G

MIRÓ Ò, CASADEMONT J, JARRETA D, GAU JM, URBANO-MARQUEZ À, CARDELLACH F, Skeletal muscle mitochondrial function in polymyalgia rheumica and in giant cell arteritis, 568

MITJAVILA F see Formiga F

MIURA M see Naito K Міулмото S see Chen JR; Takahashi M Міуовні Y see Atsumi T

MMP see Matrix metalloproteinase

Moga I see Formiga F

MOHD M see Al-Adsani A MOLA EM see Villaverde V

MOLAD Y see Krause I

MOLLNES TE, HAGA H-J, BRUN JG, NIELSEN EW, SJÖHOLM A, STURFELDT G, MARTENSSON U, BERGH K, REKVIG OP. Complement activation in patients with systemic lupus erythematosus without nephritis, 933 MONTAGNA D see Martini A

MONTEGUCCO C, CAPORALI R, ROSSI S, EPIS O. Rheumatoid arthritis in beta-thalassaemia trait, (L) 1021

MOORE GF see O'Dell JR

MORAN CJ see Stuart BM MORAND EF see Kipen Y

MORATELLI S see Trotta F

Moriyama A see Muro H

MORRIS IM. Estimating the prevalence of delayed median nerve conduction in the general population, (L) 688

MORRISON E, Capell HA. Corticosteroids in rheumatoid arthritis-the case against, (E) 97

MORTELMANS L see Blockmans D

MORTON SJ see Hardy CJ

Mosca M see Swaak AJG

Moutsopoulos HM see Mavragani CP; Swaak AJG

MUCINELLI M see Gallerani M Muir KR see Hardy CJ

MUKAMEL M see Krause I

MUKOFUJIWARA Y see Muro H

MÜLLER A see Oelzner P

MÜLLER-LADNER U, BOSSERHOFF AK, DREHER K, HEIN R, NEIDHART M, GAY S, SCHLMERICH J, BUETTNER R, LANG B. MIA (melanoma inhibitory activity): a potential serum marker for rheumatoid arthritis, 148

MUÑOZ-FERNÁNDEZ S see Villaverde V

Munro J see Brazier JE

MURAT I see Ames PRJ MURDACA G see Filaci G

Muro H, Waguri-Nagaya Y, Mukofujiwara Y, Iwahashi T, Otsuka T, MATSUI N, MORIYAMA A, ASAI K, KATO T. Autocrine induction of gliostatin/platelet-derived endothelial cell growth factor (GLS/PD-ECGF) and GLS-induced expression of matrix metalloproteinases in rheumatoid arthritis synoviocytes, 1195

MURRAY FE see Shah AA

Musculoskeletal conditions in children attending two Togolese hospitals, 1010

MYLLYKANGAS-LUOSUJÄRVI R, AHO K, KAUTIAINEN H, HAKALA M. Amyloidosis in a nationwide series of 1666 subjects with rheumatoid arthritis who died during 1989 in Finland, 499

Myositis

the multiple picture of cardiac involvement in, 1153

palpable nodular, with sarcoidosis: periostitis and large-vessel vasculitis simulating Takayasu's arteritis, (L) 287

recurrent focal (L) 1295

NAGANO A T see Naito K

NAGANO S see Nakashima H NAGASAWA K see Horiuchi T

NAITO K, TAKAHASHI M, KUSHIDA K, SUZUKI M, OHISHI T, MIURA M, INOUE T, NAGANO A. Measurement of matrix metalloproteinases (MMPs) and tissue inhibitor of metalloproteinases-1 (TIMP-1) in patients with knee osteoarthritis: comparison with generalized osteoarthritis, 510

NAITOU K see Takahashi M

NAKACHE J-P see Calin A; Dougados M

NAKAMURA K see Manabe N

NAKAMURA T see Seki K

NAKASHIMA H, AKAHOSHI M, TANAKA Y, YAMAOKA K, OGAMI E, NAGANO S, ARINOBU Y, NIIRO H, OTSUKA T, NIHO Y. POlymorphisms within the interleukin-10 receptor cDNA gene (IL10R) in Japanese patients with systemic lupus erythematosus, 1142

NASR FW see Hajj-Ali RA

NASRALLA M see Haier J
NAUGHTON MA, BATTAGLIA E, O'BRIEN S, WALPORT MJ, BOTTO M. Identification of thiopurine methyltransferase (TPMT) polymorphisms cannot predict myelosuppression in systemic lupus erythematosus patients taking azathioprine, 640

NAVARRO-BLASCO FJ. Alfuzosin is not associated with dermatomyositis, (L) 579

Neck pain, acupuncture for: a systematic review of randomized controlled trials, 143 Neidhart M see Müller-Ladner U

Nephropathy, glomerular and tubular proteinuria as markers of in arthritis, 28

Nephrotic syndrome, Adult Still's disease with, at presentation, (L) 283 NEUMANN-ANDERSEN G. Two ultraviolet radiation-induced episodes of optic

neuritis in a patient with antinuclear antibody, (L) 1296

NEURE L see Yin Z NEVEZ G, PRUNA A, JOUNIEAUX V, MAKDASSI R, TOTET A, RACCURT C. A search for Pneumocystitis carinii DNA by polymerase chain reaction on bronchoalveolar lavage fluids from patients with Wegener's granulomatosis, (L) 1025

NIAZY MN see Al-Adsani A

NICOLSON GL see Haier J

NIEDERSTADT C, HAPP T, TATSIS E, SCHNABEL A, STEINHOFF J. Glomerular and tubular proteinuria as markers of nephropathy in arthritis, 28

NIELSEN EW see Mollnes TE

NIEMAN F see van der Heijde D

NIERS TMH see van der Graaff WL

NIETHAMMER D, KÜMMERLE-DESCHNER J, DANNECKER GE. Side-effects of long-term immunosuppression versus morbidity in autologous stem cell rescue: striking the balance, 747

Nіно Y see Nakashima Н

NIIRO H see Nakashima H

Nimesulide

and diclofenac, comparative efficacy and safety in patients with acute shoulder, (S1) 33 effect of compared with naproxen on the human gastrointestinal tract,

(S1) 24

effect on glucocorticoid receptor activity in human synovial fibroblasts, (S1) 11

meta-analysis of controlled studies, (S1) 33

molecular model of the interaction with human cyclooxygenase-2, (S1) 14

overview of, (S1) 1

safety related to its pharmacokinetics: assessment of adverse reactions,

Nimesulide cyclooxygenase-2 selectivity, in vivo assessment of, (S1) 19

NINET J see Duhaut P

NISHIOKA K see Maeda T NISKIZAKA H see Horiuchi T

Nitric oxide synthesis in peripheral blood mononuclear and polymorphonuclear cells from patients with systemic sclerosis, (L) 1301

NIVISON-SMITH I see Snowden JA

NJOBVU P, HUNT I, POPE D, MACFARLANE G. Pain amongst ethnic minority groups of South Asian origin in the United Kingdom: a review, 1184 NOBLE BA see Bhakta BB

Noivo Y see Wada M

Non-steroidal anti-inflammatory drug trial, ankylosing spondylitis: what is the optimum duration of a clinical study? One year versus a six Non-steroidal anti-inflammatory drugs

back to the future. (E) 693

gastrointestinal toxicity of: the effect of nimesulide compared with naproxen, (S1) 24

NORDBORG C see Petursdottir V NORDBORG E see Petursdottir V

NORDMANN Y see Chalem P Nourooz-Zahed J see Ames PRJ

5'-Nucleotidase as a marker of both general and local inflammation in rheumatoid arthritis patients, 391

NUYTS J see Blockmans D

O'BRIEN S see Naughton MA

O'CONNOR P see McGonagle D

O'DELL JR, MOORE GF, KLASSEN LW. Combination therapy in rheumatoid arthritis: a comment, (L) 577

O'DELL JR, SCOTT DL. Rheumatoid arthritis: new developments in the use of exisitng therapies, (S2) 24

O'DUFFY EK, OLIVER FJ, CHATTERS, WALKER H, LLOYD DC, EDWARDS JCW, ELL PJ. Chromosomal analysis of peripheral lymphocytes of patients before and after radiation synovectomy with samarium-153 particulate hydroxyapatite, 316

O'GRADAIGH D, WATTS R, GLENN A, SCOTT DGI. An unusual cause of blindness in Wegener's granulomatosis, (L) 578

O'HARA M see Helliwell PS

O'SULLIVAN MM see Sinha A

ÖCAL L see Gül A

Осні Т see Tomita Т ODA H see Manabe N

OELZNER P, FRANKE S, MÜLLER A, HEIN G, STEIN G. Relationship between soluble markers of immune activation and bone turnover in postmenopausal women with rheumatoid arthritis, 841

OGAMI E see Nakashima H

Ogasawara H, Kaneko H, Hishikawa T, Sekigawa I, Takasaki Y, Hashimoto H, Hirose S, Kaneko Y, Maruyama N. Molecular mimicty between human endogenous retrovirus clone 4-1 and HLA class I antigen with reference to the pathogenesis of systemic lupus erythematosus, (L) 1163

Онізні Т see Naito K

Онта Т see Inokuma S OHZONO K see Tomita T

OKSEL F see Aksu K

Oku H see Tomita T

OKUR H see Yetgin S

OLCAY L see Yetgin S

OLIVER FJ see O'Duffy EK

OLIVIERI J see Wicki J

OLLIER WER see Donn RP; Mattey DL OLUSI SO see Al-Awadhi AM

ONIANKITAN O see Mijiyawa M

OPITZ C see Riemekasten G

Optic neuritis, two ultraviolet radiation-induced episodes of in a patient with antinuclear antibody, (L) 1296

Oral desensitization in patients with chronic tophaceous gout and allopurinol hypersensitivity, (L) 85

ORTIZ DE URBINA F see Cuende E

Osteoarthritis

determinants of WOMAC function, pain and stiffness scores: evidence for the role of low back pain, symptom counts, fatigue and depression in, 355

hip, and acetabular dysplasia prevalence in Kamitonda: from a longitudinal population-based epidemiological study of rheumatic diseases in Japan, 793

knee see Knee osteoarthritis

measurement of matrix metalloproteinases (MMPs) and tissue inhibitor of metalloproteinases-1 (TIMP-1) in patients with knee osteoarthritis: comparison with generalized, 510

and rheumatoid arthritis

comparison of the concentrations of pentosidine in the synovial fluid, serum and urine, 1275

comparison of free and peptide-bound pyridinoline cross-links excretion, 133

surfactant treatment for, (L) 1020

twins and the genetic architecture of, (E) 583

Osteoporosis, corticosteroid-induced, use of prophylaxis in hospital practice, (L) 577

ØSTERGAARD M see Klarlund M

Ota H see Inoue K Otsuka T see Muro H; Nakashima H

Outcome measurement

generic and condition-specific, for people with osteoarthritis of the

responsiveness of in juvenile chronic arthritis, 176

spontaneous ambulatory activity as a quantifiable, for rheumatoid arthritis, 1234

Outcomes and health economic issues surrounding the use of biological therapies in patients with rheumatoid arthritis, (S2) 48

Oxidative stress in systemic lupus erythematosus and allied conditions with vascular involvement, 529

ÖZCELIK T see Gül A

OZDOGAN H see Basaranoglu M

Özekmekçi S see Ertan S

OZEN S see Yetgin S

Özgül A, TANYÜKSEL M, YAZICIOĞLU K, ARPACIOĞLU O. Sacroiliitis associated with Cryptosporidium parvum in an HLA-B27-negative patient,

PAC M see Formiga F

Pain

amongst ethnic minority groups of South Asian origin in the United Kingdom: a review, 1184

chronic widespread, prevalence and associated features of in the community using the 'Manchester' definition, 275

steroid injection for: evidence of short-term effectiveness, 974

a systematic review of treatments, 968

neck, acupuncture for: a systematic review of randomized controlled trials, 143

and sleep disturbances with special reference to fibromyalgia and rheumatoid arthritis, (E) 1035

see also Low back pain PALEY MN see Clunie GPR

Palindromic morphoea, extreme hypereosinophilia in a young male patient presenting with, (L) 377

PALMER BP see Hardy CJ

PANAYI G see Emery P

PANAYI GS. Targeting of cells involved in the pathogenesis of rheumatoid arthritis, (S2) 8

Panayi GS *see also* Coakley G; Cortigall VM; Dolan AL Papo T, Le Thi Huong D, Wiederkehr JL, Woehl-Kremer B, Bletry O, Wechsler B, Godeau P, Piette JC. Etoposide in Wegener's granulomatosis, (L) 473

PARCHIN-GENESTE N see Maillefert JF

PARHAM D see Hopkinson N

PARK KEJ, CHIPPS DR, BENSON EM. Necrotizing vasculitis secondary to propylthiouracil presenting as purpura fulminans, (L) 790

PARRY E see Middleton J

PARTSCH G see Čeponis A

PASCUAL E. Management of crystal arthritis, (E) 912 PASQUIER J see Duhaut P

PASTERNAK A see Pertovaara M

PATEL S see Johnson SM PATEL SD see Hall FC

Pătiălă H see Čeponis A

PATRONO C see Brooks P

PAULE B see Clerc D

PAULUS HE, KALDEN JR. Workshop summary: the emerging role of biologicals in the treatment of patients with newly diagnosed rheumatoid arthritis, (S2) 45

PAWLOTSKY Y see Perdriger A

PAYNE L see Persey M
Pease CT, Bhakta BB, Devlin J, Emery P. Does the age of onset of rheumatoid arthritis influence phenotype?: a prospective study of outcome and prognostic factors, 228

PÉCLAT V see So A

PELLETIER J-P, DI BATTISTA JA, ZHANG M, FERNANDES J, ALAADDINE N, MARTEL-PELLETIER J. Effect of nimesulide on glucocorticoid receptor activity in human synovial fibroblasts, (S1) 11

D-Penicillamine, therapeutic response to in rheumatoid arthritis: influence of glutathione S-transferase polymorphisms, 43

Pentosidine, comparison of concentrations in the synovial fluid, serum and urine of patients with rheumatoid arthritis and osteoarthritis, 1275

PEOC'H M see Dumolard A PERDRIGER A, GUGGENBUHL P, CHALES G, YAOUANG J, QUELVENNEC E, BONNARD MN, PAWLOTSKY Y, SEMANA G. Positive association of the HLA DMB1\*0101-0101 genotype with rheumatoid arthritis, 448

PÉREZ C see Garcia-Nieto R

PERISTERAKIS A see Tatsis G

PERKOVIČ T see Tomšič M

PEROTTI C see Martini A

PERSEY M, PAYNE L, WEBLEY M. Finasteride: a steroid-unsparing drug?, (L) 285

PERTOVAARA M, KORPELA M, KOURI T, PASTERNAK A. The occurrence of

renal involvement in primary Sjögren's syndrome: a study of 78 patients, 1113

PETER HH see Jacki SH

PETTY RE. Prognosis in children with rheumatic diseases: justification for consideration of new therapies, 739

PETTY RE see also Kuis W; Wulffraat NM

PETURSDOTTIR V, JOHANSSON H, NORDBORG E, NORDBORG C. The epidemiology of biopsy-positive giant cell arteritis: special reference to cyclic fluctuations, 1208

Pharmaco-economic considerations, impact on the utilization of novel antirheumatic therapies, (S2) 41

PHELIP X see Dumolard A

PIETTE IC see Papo T

PINÈDE L see Duhaut P

PIPITONE N see Cavallo G

PIROTH C see Maillefert JF Pistorio A see Ruperto N

PITSILLIDES AA, SKERRY TM, EDWARDS JCW. Joint immobilization reduces synovial fluid hyaluronan concentration and is accompanied by changes in the synovial intimal cell populations, 1108

PITZALIS C see Gerli R

Pizzolato G see Wicki J

PLANT MJ see Mattey DL

PLUVINAGE P see Seror P

PODSCHUM R see Sahly H POKORNY G see Swaak AJG

Polyarteritis nodosa

cutaneous, with seronegative arthritis, (L) 1161

familial Mediterranean fever and, complicated by spontaneous bilateral perirenal and subcapsular splenic haemorrhage, (L) 794

related to hepatitis B virus, successful treatment of with a combination of lamivudine and interferon alpha, (L) 183

Polyarticular heterotopic ossification complicating critical illness, 1145 Polymerase chain reaction

genus-specific, and sequence anaylsis, detection of mycobacteria in joint samples from patients with arthritis using, 547

nested, broad-range, detection of DNA from a range of bacterial species in the joints of patients with a variety of arthritides using, 260

a search for Pneumocystitis carinii DNA by, on bronchoalveolar lavage fluids from patients with Wegener's granulomatosis, (L) 1025

Polymyalgia rheumatica

are pregnancies a protective factor? A prospective multicentre casecontrol study, 118

new arguments for a vasculitic nature of using positron emission tomography, 444

skeletal muscle mitochondrial function in, 568

Polymyositis, safety of a home exercise programme: a pilot study, 608 Polytopic spondylodiscitis secondary to Streptococcus bovis endocarditis, (L) 686

POPE D see Niobyu P Porphyria cutanea tarda

affecting a rheumatoid arthritis patient treated with methotrexate: association or coincidence?, 453

in a patient with systemic lupus erythematosus, (L) 1166

Positron emission tomography, new arguments for a vasculitic nature of polymyalgia rheumatica using, 444

POST WJ see Kraan MC

Postgraduate degrees for rheumatology trainees: an options appraisal of MD, PhD and MSc degrees, 1290
POSTHUMUS MD, LIMBURG PC, WESTRA J, CATS HA, STEWART RE, VAN

LEEUWEN MA, VAN RUSWUK MH. Serum levels of matrix metalloproteinase-3 in relation to the development of radiological damage in patients with early rheumatoid arthritis, 1081

POULTON KV see Mattey DL POWELL RJ see Hardy CJ

PRABHA K see Al-Awadhi AM

Pregnancy

-induced flare in systemic lupus erythematosus and remission in rheumatoid arthritis, 496 giant cell arteritis and polymyalgia rheumatica: a protective factor? A

prospective, multicentre case-control study, 118 intravenous immunoglobulins for Adult Still's disease and, (L) 1024 outcome and family size in systemic lupus erythmatosus: a case-control

study, 559 Pregnancy loss, recurrent, and autoantibody profile in autoimmune diseases, 1228

PRESTON A see Fraser A

PREVOO MLL see Albers JMC

PRICE PA see Johansen JS

Primary care rheumatology-leading the way?, (E) 1174

Primary care-led rheumatology services, comparison with secondary care provision, (L) 1294

PRINS APA see van der Graaff WL

PRITCHARD MH, PUGH N, WRIGHT I, BROWNLEE M. A vascular basis for repetitive strain injury, 636

PRITCHARD T see Ferry S PRIULI GB see Mijiyawa M

Probucol improves symptoms and reduces lipoprotein oxidation susceptibility in patients with Raynaud's phenomenon, 309

Prolactin, the immune response and lupus, (E) 296

Propylthiouracil, necrotizing vasculitis secondary to, presenting as purpura fulminans, (L) 790

PROTHEROE A, EDWARDS JCW, SIMMONS A, MACLENNAN K, SELBY P. Remission of inflammatory arthropathy in association with anti-CD20 therapy for non-Hodgkin's lymphoma, 1150

PROUDMAN S see Emery I

PROUSE P see Solymossy C

PRUNA A see Nevez G

Psoriatic arthritis

associated with chondrolysis and osteolysis, (L) 86

clinical, radiographic and HLA associations as markers for different patterns of, 332

comparative quantitative morphometric study of cell apoptosis in synovial membranes in, reactive and rheumatoid arthritis, 431 imaging of pelvic fracture, (L) 902

or SAPHO syndrome? A familial case study, 463

transmission of by allogeneic bone marrow transplantation for chronic myelogenous leukaemia from an HLA-identical donor, (L) 89

PUGH N see Pritchard MH Pujol R see Formiga F

PUOLAKKAINEN M see Hannu T

PUPPO F see Filaci G

Purpura fulminans, necrotizing vasculitis secondary to propylthiouracil presenting as, (L) 790

Pyridinium cross-links, urinary, disease activity and disease subsets, relationship in ankylosing spondylitis, 21

Pyridinoline cross-links excretion, free and peptide-bound, comparison of in rheumatoid arthritis and osteoarthritis, 133

QUELVENNEC E see Perdriger A

OUILLARD J see Clerc D

QVINDESLAND A, JÓNSSON H. Articular hypermobility in Icelandic 12-yearolds, 1014

RACCURT C see Nevez G

RADBRUCH A see Yin Z

Radiation synovectomy with samarium-153 particulate hydroxyapatite. Chromosomal analysis of peripheral lymphocytes of patients before and after, 316

Radiographic changes, and quality of life in early rheumatoid arthritis. A 12-month randomized controlled trial of patient education on, 303

Radiographic, clinical and HLA associations as markers for different patterns of psoriatic arthritis, 332

Radiographic joint space in rheumatoid acromioclavicular joints: a 15 year prospective follow-up study in 74 patients, 1104

Radiographs, reading in chronological order, in pairs or as single films has important implications for the discriminative power of rheumatoid arthritis clinical trials, 1213

Radiological assessment in rheumatoid arthritis, Sharp/van der Heijde, reliability and sensitivity to change of a simplification of, 941

Radiological damage in patients with early rheumatoid arthritis, serum levels of matrix metalloproteinase-3 in relation to the development of. 1081

Radiological outcome in rheumatoid arthritis, influence of HLA-DRBI alleles encoding the DERAA amino acid motif on, 1221

RADSTAKE TRDJ see Barrera P

RADWAY-BRIGHT EL, INANC M, ISENBERG DA. Animal models of the antiphospholipid syndrome, 591

RAFTERY MJ see Smith PR

RAINSFORD KD. Relationship of nimesulide safety to its pharmacokinetics: assessment of adverse reactions, (S1) 4

RAMAGE JM, GASTON JSH. Depressed proliferative responses by peripheral blood mononuclear cells from early arthritis patients to mycobacterial heat shock protein 60, 631

RAMAGE P see Dawson J

RASKER H see Armstrong R

RASPE H see Dieppe P RAU R see van der Heijde D

RAVELLI A see Martini A; Ruperto N

Raynaud's phenomenon

primary, case-control study of candidate vasoactive meditator genes, 1094

probucol improves symptoms and reduces lipoprotein oxidation susceptibility in, 309

REKVIG OP see Molines TE

Renal failure, normotensive, in a patient with systemic sclerosis and p-antineutrophil cytoplasmic autoantibodies which developed into Paget's disease of bone after immunosuppressive therapy, (L) 190

Renal function, irreversible loss of, cyclosporin A therapy in rheumatoid arthritis: only strict application of the guidelines for safe use can prevent, 254

Renal involvement, occurrence of in primary Sjögren's syndrome: a study of 78 patients, 1113

Repetitive strain injury, a vascular basis for, 636

Reproductive failure prior to the onset of clinical autoimmune disease, (E) 485

Rheumatoid arthritis

50 years of glucocorticoid treatment in, (E) 100

allergic pancytopenia to trimethoprim-sulphamethoxazole for Pneumocystis carinii pneumonia following methotrexate treatment for, (L) 475

amyloidosis in a nationwide series of 1666 subjects with, who died during 1989 in Finland, 499

angio-neurotic oedema associated with methotrexate treatment. 908 anticardiolipin antibodies, free protein S levels and thrombosis: a survey in a selected population of patients, 675

apocynin, a plant-derived, cartilage-saving drug, might be useful in treatment, 1088

are the risks of haemopoietic stem cell transplantation acceptable, 321 assessing the role of anti-tumour necrosis factor therapy in the treatment of, recapitulation of the round-table discussion, (S2) 50

assessment of disease activity using magnetic resonance imaging; quantification of pannus volume in the hands, (L) 680

autocrine induction of gliostatin/platelet-derived endothelial cell growth factor (GLS/PD-ECGF) and GLS-induced expression of matrix metalloproteinases in synoviocytes, 1195

in beta-thalassaemia trait, (L) 1021

CD5 and CD23 expression on B cells in peripheral blood and synovial fluid: relationship with interleukin-4, soluble CD23 and tumour necrosis factor alpha levels, 325

changes in plasma free fatty acid concentrations in patients during fasting and their effects upon T-lymphocyte proliferation, 948

clinical trials, reading radiographs in chronological order, in pairs or as single films has important implications for the discriminative power of, 1213

combination therapy (L) 577, 789 comparative quantitative morphometric study of cell apoptosis in synovial membranes in psoriatic, reactive and, 431

corticosteroids in-the case against, (E) 97

cyclosporin A therapy in: only strict application of the guidelines for safe use can prevent irreversible renal function loss, 254

cytokine production by synovial T cells in, 202 and depression, effects of dothiepin on subjects with, 959

detection of mycoplasmal infections in blood of patients, 504

determinants of WOMAC function, pain and stiffness scores: evidence for the role of low back pain, symptom counts, fatigue and depression in, 355

diet therapy for the patient with?, (E) 1039

does the age of onset influence phenotype?: a prospective study of outcome and prognostic factors, 228

12-month randomized controlled trial of patient education on radiographic changes and quality of life, 303

the case for corticosteroids in the treatment of, (E) 95

changes in serum chondroitin sulphate epitopes 3-B-3 and 7-D-4, 837 preliminary study of ultrasound aspiration of bone erosion, 329 serum levels of matrix metalloproteinase-3 in relation to the development of radiological damage in, 1081

soluble CD30 in as a predictor of good response to second-line therapy, 1282

effect of acupuncture on: a randomized, placebo-controlled crossover study, 864

effect of methotrexate on blood purine and pyrimidine levels in, 997 elevated ratio of interferon gamma-/interleukin-4-positive T cells found in synovial fluid and synovial membrane of patients can be changed by interleukin-4 but not by interleukin-10 or transforming growth factor beta, 1058

establishment of nurse-like stromal cells from bone marrow of patients: indication of characteristic bone marrow microenvironment, 854

evaluation of a new Apo-1/Fas promoter polymorphism in, 645

extra-articular: prevalence and mortality, 668

extrapyramidal type rigidity in, 627

familial aggregation in The Netherlands: a cross-sectional hospitalbased survey, 415

female patients, relationships between serum 17-β-oestradiol and anticardiolipin antibody concentrations in, (L) 1159

finger joint synovitis in: quantitative assessment by magnetic resonance imaging, 66

future trends in the management of, (S2) 1

future trends in the treatment of: cytokine targets, (S2) 11

gene transfer as therapy for: why, what and how?, (E) 804 glucosteroids in the management of, 6

health outcomes and health economic issues surrounding the use of biological therapies in, (S2) 48

high-dose chemotherapy in: a report of five cases, (L) 181

how may quality of life be enhanced by current and future treat-ments?, (S2) 35

increased Ed-B fibronectin plasma levels in spondyloarthropathies: comparison with RA patients and a healthy population, 1099 influence of HLA-DRB1 alleles encoding the DERAA amino acid motif on radiological outcome in, 1221

influence of a partially HLA-matched blood transfusion on the disease activity of, 53

interleukin-10 as an explanation for pregnancy-induced remission in 496

involvement of fibroblast growth factor-2 in joint destruction, 714 is there a defect in cortisol production?, (E) 298

new developments in the use of exisitng therapies, (\$2) 24 newly diagnosed

emerging role of biologicals in the treatment of, (S2) 45

management of patients with, (S2) 27

5'-nucleotidase as a marker of both general and local inflammation in 391

oral carriage of staphylococci in, 572

oral low-dose glucocorticosteroids as compared with intravenous methylprednisolone pulses in the treatment of, (L) 1304 and osteoarthritis

comparison of the concentrations of pentosidine in the synovial fluid, serum and urine, 1275

comparison of free and peptide-bound pyridinoline cross-links excretion, 133

pain and sleep disturbances with special reference to, (E) 1035 peripheral ulcerative keratitis 'corneal melt' and: a case series, 1245 porphyria cutanea tarda affecting a patient treated with methotrexate: association or coincidence?, 453

positive association of the HLA DMB1\*0101-0101 genotype with,

post-menopausal women, relationship between soluble markers of immune activation and bone turnover in, 841 predicting normal 'grip' strength for patients, 521

prevalence of hypothyroidism among Arabs with, (L) 472

management of, (S2) 32

role of biologicals in the treatment of, (S2) 47

relative proportions of secreted interleukin-2 and interleukin-10 determine the magnitude of T-cell proliferation to the recall antigen turberculin purified protein derivative, 1203

relevance of tumour necrosis factor alpha gene polymorphism in, (L) 681

reliability and sensitivity to change of a simplification of the Sharp/ van der Heijde radiological assessment in, 941

role of current strategies in the future treatment of, (S2) 19

role of cytokines in the pathogenesis of, (S2) 3

sensitivity and specificity of anti-Sa autoantibodies for, 155

seropositive, work disability in an cohort of patients with: a 20 year study, 1138

serum lysozyme: a potential marker of monocyte/macrophage activity in 1249

serum MMP-3 in: correlation with systemic inflammation but not with erosive status, 407

serum YKL-40 concentrations in: relation to disease activity, 618 silastic prostheses-a forgotten cause of lymphadenopathy in, (L) 480 sleep apnoea caused by, (L) 679, 680

socio-economic consequences in the first years of the disease, 423 spontaneous ambulatory activity as a quantifiable outcome measure for, 1234

stress system response and: a multilevel approach, 1050 targeted therapies in: the need for action, (E) 911

targeting of cells involved in the pathogenesis of, (S2) 8

Th1-type cytokine mRNA in mononuclear cells induced by streptococcal pyrogenic exotoxin A, (L) 1022

therapeutic response to D-penicillamine in: influence of glutathione S-transferase polymorphisms, 43

RIBERA A see Marsal S

RICARD-BLUM S see Toussirot E

RICHARDS PJ, AMOS N, WILLIAMS AS, WILLIAMS BD. Pro-inflammatory effects of the aminobisphosphonate ibandronate in vitro and in vivo, 984 RICHARDS PJ, WILLIAMS AS, GOODFELLOW RM, WILLIAMS BD. Liposomal clodronate eliminates synovial macrophages, reduces inflammation and ameliorates joint destruction in antigen-induced arthritis, 818

RICHART C see Garcia-Pardo G

RIEMEKASTEN G, OPITZ C, AUDRING H, BARTHELMES H, MEYER R, HIEPE F, BURMESTER GR. Beware of the heart: the multiple picture of cardiac involvement in myositis. 1153

RIGBY SP see Schott JM

RIVERA A see Formiga F ROBERTS Z see Denton CP

ROBINZON B, CUTOLO M. Should dehydroepiandrosterone replacement therapy be provided with glucocorticoids? 488

ROBSON T see Szczepanska K

ROLF C, ENGSTRÖM B, BEAUCHARD C, JACOBS LD, LE LIBOUX A. Intraarticular absorption and distribution of ketoprofen after topical plaster application and oral intake in 100 patients undergoing knee arthrosсору, 564

ROLLO L see Cavallo G

ROONEY B see McNally R ROPOLO F see Filaci G

ROY DK, GEORGE A, CHATTOPADHYAY C, GRENNAN DM. Splenic infarction in a patient with Wegener's granulomatosis, (L) 1162

ROZMAN B see Swaak AJG; Tomšič M

RUDWALEIT M see Brandt J

RUOF J, BRÜHLMANN P, MICHEL BA, STUCKI G. Development and validation of a self-administered systemic sclerosis questionnaire (SySQ), 535

RUPERTO N, RAVELLI A, FALCINI F, LEPORE L, BUONCOMPAGNI A, GERLONI V, BARDARE M, CORTIS E, ZULIAN F, SARDELLA ML, GIOVANNI STRANO C, ALESSIO M, ALPIGIANI MG, MIGLIAVACCA D, PISTORIO A, VIOLA S, MARTINI A. Responsiveness of outcome measurements in juvenile chronic arthritis, 176 RUSTAN AC see Fraser DA

RYAN PJ see Seeman E

RYZEWSKA A see Szczepanska K

SAATCI U see Yetgin S

SABADINI L see Cavallo G

Sacral insufficiency fractures, an unsuspected cause of low back pain, (L) 90. 903

Sacroiliitis

acute anterior uveitis in a patient with, and acne conglobata, (L) 797 associated with Cryptosporidium parvum in an HLA-B27-negative patient, (L) 288

studying patients with inflammatory back pain and arthritis of the lower limbs clinically and by magnetic resonance imaging: many, but not all, patients with, have spondyloarthropathy, 831

SAHLY H, PODSCHUN R, ULLMAN U. Increased antibody responses to Klebsiella serotypes K26, K36 and K50 in patients with ankylosing spondylitis, (L) 481

SAINIO P see Mäkelä M

SAINTE-LAUDY J see Chevrel G

SALMON M, GORDON C. The role of apoptosis in systemic lupus erythematosus, 1177

SALMON M see also Gordon C

SALTI IA see Hajj-Ali RA Samarium-153 particulate hydroxyapatite

chromosomal analysis of peripheral lymphocytes of patients before and after radiation synovectomy with, 316

intra-articular, changes in articular synovial lining volume measured by magnetic resonance in a randomized, double-blind, controlled trial of for chronic knee synovitis, 113

SANCHEZ-HARGUINDEY L see Fernández-Gutiérrez B

SÁNCHEZ-IBARROLA A see Loza E

SANCHEZ-RUIS A see Jimenez-Mazuecos J

SAPHO syndrome or psoriatic arthritis? A familial case study, 463

SARASIN F see Wicki J

Sarcoidosis with palpable nodular myositis: periositis and large-vessel vasculitis simulating Takayasu's arteritis, (L) 287

SARDELLA ML see Ruperto N

SARI-KOUZEL H, HERRICK AL, FREEMONT AJ, MARCUSON RW, JAYSON MI. Giant cell arteritis in a patient with limited cutaneous systemic sclerosis, (L) 479

SATHANATHAN R see David J

SATO S see Hasegawa M SAYINER A see Aksu K

SCHATTENKIRCHNER M see Anders H-J SCHELLEKENS JFP see Blaauw AAM

SCHNABEL A see Niederstadt C

SCHNEIDER M see Swaak AJG

SCHLMERICH J see Müller-Ladner U

SCHOTT JM, RIGBY SP, McNALLY JD, KEAT A, HIGGENS CS. Oral methotrexate: the hazard of different tablet strengths, (L) 382

SCHOULS LM see van der Heijden IM

SCHREUDER GMTh see van der Horst-Bruinsma

SCHULMAN S see Barzilai A

Scleroderma see Systemic sclerosis

SCOTT DL. Changing times, (E) 1 SCOTT DL see also Houssien DA; O'Dell JR

SCUDELETTI M see Filaci G

SEEDHOM BB see Shepherd DET: Yao JO

SEEMAN E, RYAN PJ. Bone turnover and bone mass in polymyalgia rheumatica, (L) 286

SEKI K, TAKAHASHI R, NAKAMURA T. Psoriatic arthritis associated with chondrolysis and osteolysis, (L) 86

Sekigawa I see Ogasawara H

SELBY P see Protheroe A

SEMANA G see Perdriger A

SERIOLO B, ACCARDO S, GARNERO A, FASCIOLO D, CUTOLO M. Anticardiolipin antibodies, free protein S levels and thrombosis: a survey in a selected population of rheumatoid arthritis patients, 675

SERIOLO B, CUTOLO M, GARNERO A, ACCARDO S. Relationships between serum 17-β-oestradiol and anticardiolipin antibody concentrations in female patients with rheumatoid arthritis, (L) 1159

Seronegative oligoarthritis in the course of refractory anaemia with excess blasts, (L) 373

SEROR P, PLUVINAGE P, LECOQ d'ANDRE F, BENAMOU P, ATTUIL G. Frequency of sepsis after local corticosteroid injection (an inquiry on 1 160 000 injections in rheumatological private practice in France), 1272

Setoguchi K see Inokuma S

SEVERENS JL see Albers JMC SEYDOUX D see Duhaut P

SHAH AA, MURRAY FE, FITZGERALD DJ. The in vivo assessment of nimesulide cyclooxygenase-2 selectivity, (S1) 19

SHAHDAND S see Fadilah SAW

SHANAHAN EM, HUTCHINSON M, HANLEY SD, BRESNIHAN B. Giant cell arteritis presenting as lateral medullary syndrome, (L.) 188 Shanahan EM, Sheahan K, McDonald K, Bresnihan B. Dilated cardi-

omyopathy in Wegener's granulomatosis, (L) 1164 Sheahan K see Shanahan EM

SHEPHERD DET, SEEDHOM BB. The 'instantaneous' compressive modulus of human articular cartilage in joints of the lower limb, 124

SHICHIKAWA K see Inoue K

SHIPTON EA see Dessein PH

Shoulder, acute, comparative efficacy and safety of nimesulide and diclo-fenac, (S1) 33

Shoulder complaints, long-term course of: a prospective study in general practice, 160

Shoulder girdle, Gorham-Stout disease (phantom bone) of, (L) 904

Shoulder joint impairment among Finns aged 30 years or over: prevalence, risk factors and co-morbidity, 656

SIBILIA J see Maillefert JF

SIEGERT S see Yin Z

SIEPER J see Brandt J; Wilkinson NZ; Yin Z

Silastic prostheses-a forgotten cause of lymphadenopathy in rheumatoid arthritis, (L) 480

SILMAN AJ see Bhakta BB; Ferry S; Hunt IM; Thomas E

SIMMONDS HA see Smolenska Z SIMMONS A see Protheroe A

SINGH BB see Berman BM

SINHA A, DIXON N, O'SULLIVAN MM, SOWDEN JM. Porphyria cutanea tarda in a patient with systemic lupus erythematosus, (L) 1166

Siögren's syndrome

and autoimmune myelofibrosis: an uncommon association, (L) 370 a community-based study of prevelance and impact-comment on the article by Thomas et al, (L) 685

(L) (Reply) 686

microvascular abnormalities in: nailfold capillaroscopy, 826

prevalence in Slovenia, 164

primary

in the North East of England: a long-term follow-up study, 245 occurrence of renal involvement in: a study of 78 patients, 1113 zidovudine in, 814

SJÖHOLM A see Mollnes TE

SKERRY TM see Pitsillides AA

SKINNER CJ see Smith PR

SKOPOULI FN see Tektonidou M

SLAPER-CORTENBACH ICM, WIJNGAARDEN-DU BOIS MJGJ, de VRIES-VAN ROSSEN A, BORST HPE, VAN DER LELIE H, VAN HEUGTEN HG, VERDONCK LF, WULFFRAAT NM, HOOGERBRUGGE PM. The depletion of T cells from haematopoietic stem cell transplants, 751

Sleep apnoea caused by rheumatoid arthritis, (L) 679, 680

Sleep disturbances and pain with special reference to fibromyalgia and rheumatoid arthritis, (E) 1035

Small intestinal perforation complicating Wegeners' granulomatosis, (L) 289 SMEENK RJT see Swaak AJG

SMEETS TIM see Tak PP

SMITH MD. Assessment of disease activity in rheumatoid arthritis using magnetic resonance imaging: quantification of pannus volume in the hands, (L) 680

SMITH PR. D'CRUZ D, RAFTERY MJ, DODD SM, HELBERT M, SKINNER CJ. Drug-induced lupus nephritis in HIV infection, (L) 1017

SMOLEN JS see Brooks P; Furst DE; Hueber W; Steiner G; Swaak AJG; Weisman

SMOLENSKA Z, KAZNOWSKA Z, ZARÓWNY D, SIMMONDS HA, SMOLENSKI RT. Effect of methotrexate on blood purine and pyrimidine levels in patients with rheumatoid arthritis 997

SMOLENSKI RT see Smolenska Z.

SMYTH AE, HUGHES AE, BRUCE IN, BELL AL. A case-control study of canditate vasoactive meditator genes in primary Raynaud's phenomenon, 1094

SNAITH ML see Brazier JE

SNOWDEN JA, NIVISON-SMITH I, BIGGS JC, BROOKS PM. Risk taking in patients with rheumatoid arthritis: are the risks of haemopoietic stem cell transplantation acceptable, 321
So A, Chamot AM, Péclat V, Gerster J-C. Serum MMP-3 in rheumatoid

arthritis: correlation with systemic inflammation but not with erosive status, 407

SØNDERSTRUP G see Hall FC

SOBEL JS see Winters JC

SOLYMOSSY C, DIXEY J, UTLEY M, GALLIVAN S, YOUNG A, COX N, DAVIES P, EMERY P, GOUGH A, JAMES D, PROUSE P, WILLIAMS P, WINFIELD J. Larsen scoring of digitalized X-ray images, 1127

SOWDEN JM see Sinha A SPECTOR TD see Lloyd ME; MacGregor AJ

Spinal kyphosis, biochemical analysis of posture in patients with due to ankylosing spondylitis: a pilot study, 441
Spinal manipulative procedures, forces measured during, in two age

groups, 267

Splenectomy for refractory thrombocytopenia in the antiphospholipid syndrome, 848

Splenic infarction in a patient with Wegener's granulomatosis, (L) 1162 Spondylarthropathies

concordance between abdominal scintigraphy using technetium-99m hexamethylpropylene amine oxime-labelled leucocytes and ileocolonoscopy in patients without clinical evidence of inflammatory bowel disease, 543

studying patients with inflammatory back pain and arthritis of the lower limbs clinically and by magnetic resonance imaging: many, but not all, patients with sacroiliitis have, 831

SQUIRRELL DM, WINFIELD J, Amos RS. Peripheral ulcerative keratitis 'corneal melt' and rheumatoid arthritis: a case series, 1245

SRINIVASAN U, COUGHLAN RJ. Small intestinal perforation complicating Wegeners' granulomatosis, (L) 289

Staphylococci, oral carriage of in rheumatoid arthritis, 572

STEIN G see Oelzner P

STEINER G, TOHIDAST-AKRAD M, WITZMANN G, VESELY M, STUDNICKA-BENKE A, GAL A, KUNAVER M, ZENZ P, SMOLEN JS. Cytokine production by synovial T cells in rheumatoid arthritis, 202

STEINER G see also Hueber W STEINFELD SD, DEMOLS P, VAN VOOREN J-P, COGAN E, APPELBOOM T. Zidovudine in primary Sjögren's syndrome, 814

STEINHOFF J see Niederstadt C

Stem cell transplantation

autologous

marked and sustained improvement 2 years after, in a girl with systemic sclerosis, 773

rationale for high-dose cyclophosphamide and medium-dose total body irradiation in the conditioning of children with progressive systemic and polyarticular juvenile chronic arthritis before, 762

in refractory juvenile chronic arthritis 737, 764

proposed guidelines, 777

side-effects of long-term immunosuppression versus morbidity in: striking the balance, 747

haematopoietic/haematological

for active systemic lupus erythematosus, 767

depletion of T cells from, 751

in rheumatoid arthritis: are the risks acceptable?, 321

in the treatment of severe autoimmune diseases: first experiences from an international project, 774

STENSTRÖM CH see Alexanderson H

Steroid injection for heel pain: evidence of short-term effectiveness, 974

Steroid therapy, chronic, osteopenia in young hypogonadal women with systemic lupus erythematosus receiving: a randomized controlled trial comparing calcitriol and hormone replacement therapy, 1239 Steroid trial, ARC, and its effect on clinical practice, (L) 905

Steroid-induced osteoporosis, management of: what is the current state of play?, (E) 198

STEWART O see McNally R

STEWART RE see Posthumus MD

STOLTENBERG M see Johansen JS STRANGE RC see Layton MA

STRAUSS BJG see Kipen Y

Stress system response and rheumatoid arthritis: a multilevel approach, 1050 STROOBANTS S see Blockmans D

STRUTHERS GR see Gillot TI

STUART BM, MORAN CJ. High-dose chemotherapy in rheumatoid arthritis: a report of five cases, (L) 181

STUCKI G see Houssien DA; Ruof J

STUDNICKA-BENKE A see Steiner G

STURFELDT G see Mollnes TE

STURROCK R see Emery P; Hamilton J; Jackson MS

SUGAMOTO K see Tomita T

SUGIMOTO H, TAKEDA A. Assessment of disease activity in rheumatoid arthritis using magnetic resonance imaging: quantification of pannus volume in the hands, (L) (Reply) 680

SULLI A see Filaci G.

SULTAN SM, IOANNOU Y, ISENBERG DA. A review of gastrointestinal manifestations of systemic lupus erythematosus, 917

SUMIDA T see Maeda T

Sun L-Y see Trotta F

Surfactant treatment for osteoarthritis, (L) 1020

SUTCLIFFE N, CLARKE AE, GORDON C, FAREWELL V, ISENBERG DA. The association of socio-economic status, race, psychosocial factors and outcome in patients with systemic lupus erythematosus, 1130

SUZUKI M see Chen JR; Naito K; Takahashi M

SUZUKI R see Tomita T SWAAK AJG, VAN DEN BRINK HG, SMEENK RJT, MANGER K, KALDEN JR, TOSI S, MARCHESONI A, DOMLJAN Z, ROZMAN B, LOGAR D, POKORNY G, KOVACS L, KOVACS A, VLACHOYIANNOPOULOS PG, MOUTSOPOULOS HM, CHWALINSKA-SADOWSKA H, DRATWIANKA B, KISS E, CIKES N, BRANIMIR A, SCHNEIDER M, FISCHER R, BOMBARDIERI S, MOSCA M, GRANINGER W, SMOLEN JS. Systemic lupus erythematosus: clinical features in patients with a disease duration of over 10 years, first evaluation, 953

Synovial fibroblasts, human, effect of nimesulide on glucocorticoid receptor activity in, (S1) 11

Synovial fluids, type IX collagen immunoreactive peptides in from arthritis patients, 338

Synovial intimal cell populations, joint immobilization reduces synovial fluid hyaluronan concentration and is accompanied by changes in, 1108

Synovial lining, articular, volume changes measured by magnetic resonance in a randomized, double-blind, controlled trial of intra-articular samarium-153 particulate hydroxyapatite for chronic knee synovitis, 113

Synovial membranes

collagenase, cathepsin B and cathepsin L gene expression in the, of patients with early inflammatory arthritis, 34 comparative quantitative morphometric study of cell apoptosis in, in

psoriatic, reactive and rheumatoid arthritis, 431 Synovial tissue, immunohistological analysis for differential diagnosis in

early arthritis, 1074

Synovium, rheumatoid, decreased TCR AV24AJ18+ double-negative T cells in. (L) 186

Systemic lupus erythematosus

active, haematopoietic stem cell transplantation for, 767

association of Fas/APO-1 gene polymorphism in Japanese patients, 516 association of optic neuropathy with transverse myelitis in, (L) 191 association of socio-economic status, race, psychosocial factors and

outcome in, 1130 clinical features in patients with a disease duration of over 10 years,

first evaluation, 953

complement activation in patients without nephritis, 933

complicated by haemorrhagic cystitis, does parenteral oestrogen therapy flare up disease activity?, (L) 372

evaluation of a new Apo-1/Fas promoter polymorphism in, 645 gastrointestinal manifestations of: a review, 917

high disease activity at baseline does not prevent remission, 724 identification of thiopurine methyltransferase (TPMT) polymorphisms cannot predict myelosuppression in patients taking azathioprine,

interleukin-10 as an explanation for pregnancy-induced flare 496 long-lasting remission and successful treatment of acquired factor VIII inhibitors using cyclophosphamide, 1007

molecular mimicry between human endogenous retrovirus clone 4-1 and HLA class I antigen with reference to the pathogenesis of, (L) 1163

osteopenia in young hypogonadal women with, receiving chronic steroid

therapy: a randomized controlled trial comparing calcitriol and hormone replacement therapy, 1239

oxidative stress in, and allied conditions with vascular involvement, 529 polymorphisms within the interleukin-10 receptor cDNA gene (IL10R) in Japanese patients, 1142

porphyria cutanea tarda in a patient with, (L) 1166

pregnancy outcome and family size in: a case-control study, 559

role of apoptosis in, 1177

serum leucine aminopeptidase as an activity indicator in: a study of 46 consecutive cases, 705

three year follow-up of body composition changes in pre-menstrual women with, 59 see also Lupus

Systemic sclerosis (scleroderma)

bone mineral density in, 371

case that developed under dexfenfluramine use, (L) 379

causes of mortality in, (L) 283

clinical and laboratory manifestations of in Black South Africans, 397 cyclosporin A and iloprost treatment: clinical results and interleukin-6 serum changes after 12 months of therapy, 992

enhanced production of interleukin-6 (IL-6), oncostatin M and soluble IL-6 receptor by cultured peripheral blood mononuclear cells from patients with, 612

lymphatic microangiopathy of the skin in, 221

marked and sustained improvement 2 years after autologous stem cell transplantation in a girl with, 773

nitric oxide synthesis in peripheral blood mononuclear and polymorphonuclear cells from patients with, (L) 1301

nitric oxide synthesis in peripheral blood mononuclear and polymorphonuclear cells, (L) 1301 normotensive renal failure in a patient with, and p-antineutrophil

cytoplasmic autoantibodies which developed into Paget's disease of bone after immunosuppressive therapy, (L) 190

self-administered questionnaire (SySQ), development and validation

SZCZEPANSKA K, CHRZANOWSKA-LIGHTOWLERS Z, DEGNAN BA, DIAMOND AG, ROBSON T, RYZEWSKA A, GOODACRE JA. Thl-type cytokine mRNA in rheumatoid arthritis mononuclear cells induced by streptococcal pyrogenic exotoxin A, (L) 1022

SZEBENYI B see Kirwan JR

decreased TCR AV24AJ18\* double-negative in rheumatoid synovium,

depletion of autografts for autoimmune diseases, role of, 755

depletion of from haematopoietic stem cell transplants, 751 interferon gamma- and interleukin-4-producing, quantification of in synovial fluid and peripheral blood of arthritis patients, 214

interferon gamma-/interleukin-4-positive, the elevated ratio of found in synovial fluid and synovial membrane of rheumatoid arthritis patients can be changed by interleukin-4 but not by interleukin-10 or transforming growth factor beta, 1058

synovial, cytokine production by in rheumatoid arthritis, 202

T-lymphocyte proliferation. Changes in plasma free fatty acid concentrations in rheumatoid arthritis patients during fasting and their effects upon, 948 't HART BA see Tak PP

TABAK F see Basaranoglu M

TAGER RE, TIKLY M. Clinical and laboratory manifestations of systemic

sclerosis (scleroderma) in Black South Africans, 397
TAK PP, 't HART BA, KRAAN MC, JONKER M, SMEETS TJM, BREEDVELD FC. The effects of interferon beta treatment on arthritis, 362

TAK PP see also Kraan MC

TAKAHASHI M, SUZUKI M, NAITOU K, MIYAMOTO S, KUSHIDA K. Comparison of free and peptide-bound pyridinoline cross-links excretion in rheumatoid arthritis and osteoarthritis, 133

TAKAHASHI M see also Chen JR; Naito K TAKAHASHI R see Seki K TAKANO H see Tomita T TAKASAKI Y see Ogasawara H TAKEDA A see Sugimoto H TAKEHARA K see Hasegawa M TAKEUCHI E see Tomita T TAL N see Korkmaz C TAMULAITIENÈ M see Čeponis A TANAKA Y see Nakashima H

TANIGUCHI M see Maeda T TANYÜKSEL M see Özgül A

Tassiopoulos T see Tatsis G TATSIS E see Niederstadt C

TATSIS G, VAIOPOULOS G, TASSIOPOULOS T, VOTSIOU A, PERISTERAKIS A, JORDANOGLOU J. Lung function in Adamantiades-Behçet disease, (L) TATSUO H see Wada M

TAVERNIER C see Maillefert JF

Tay C-H. Eosinophilic arthritis, 1188

TEKINAY T see Gül A

Текои A see Mijiyawa M

TEKTONIDOU M, KASKANI E, SKOPOULI FN, MOUTSOPOULOS HM. Microvascular abnormalities in Sjögren's syndrome: nailfold capillarosсору, 826

Templado JA see Cuende E Temporal arteritis, temporal <sup>67</sup>gallium uptake is increased in, 709

Ten Cate R see Vossen JM
Tenant MJ see Hamuryudan V

TENCH CM. An insight into rheumatology resources available on the World Wide Web. Reply to letters by McNally et al and Armstrong et al,

THERON J see Bonnotte B

Thiopurine methyltransferase (TPMT) polymorphisms, identification of cannot predict myelosuppression in systemic lupus erythematosus patients taking azathioprine, 640

THIODLEIFSSON B see Biarnason I

THOEN J see Fraser DA

THOMAS E, HAY EM, HAJEER A, SILMAN AJ. Sjögren's syndrome, (L) (Reply) 686

THOMAS R see Vecchio P
THOMPSON PW see Dechow E

THOMSON W see Donn RP; Mattey DL TIKLY M see Tager RE

TILLY H see Marie I

TINTURÉ T see Loza E

TOBIAS JH. Management of steroid-induced osteoporosis: what is the current state of play?, (E) 198

TOHIDAST-AKRAD M see Steiner G TOMASSINI C see Gerli R

TOMENSON B see Ash G

TOMINAGA K see Matsui K

TOMITA T, TAKEUCHI E, TOYOSAKI-MAEDA T, OKU H, KANEKO M, TAKANO H, SUGAMOTO K, OHZONO K, SUZUKI R, OCHI T. Establishment of nurse-like stromal cells from bone marrow of patients with rheumatoid arthritis: indication of characteristic bone marrow microenvironment in patients with rheumatoid arthritis, 854

TOMŠIČ M, LOGAR D, GRMEK M, PERKOVIČ T, KVEDER T. Prevalence of Sjögren's syndrome in Slovenia, 164

Tomšič M, Rozman B. Sjögren's syndrome: a community-based study of prevelance and impact-comment on the article by Thomas et al, (L) 685 Tomson TAS see van der Horst-Bruinsma

TOPALOĞLU R see Yetgin S

TORRABADELLA S see Cuende E

TORRE A see Villaverde V

TORSTEINSDÓTTIR I. HÁKANSSON L. HÄLLGREN R. GUDBJÖRNSSON B. ARVIDSON N-G, VENGE P. Serum lysozyme: a potential marker of monocyte/macrophage activity in rheumatoid arthritis, 1249

Tosi S see Swaak AJG

TOTET A see Nevez G

TOUSSAINT HM see Bot SDM

TOUSSIROT E, RICARD-BLUM S, DUMOULIN G, CEDOZ JP, WENDLING D. Relationship between urinary pyridinium cross-links, disease activity and disease subsets of ankylosing spondylitis, 21

Toussirot E see also Maillefert JF

TOWNSEND S see David J

Toyosaki-Maeda T see Tomita T

TRAYNOR A, BURT RK. Haematopoietic stem cell transplantation for active systemic lupus erythematosus, 767

TROTTA F, BAJOCCHI G, LA CORTE R, MORATELLI S, SUN L-Y. Long-lasting remission and successful treatment of acquired factor VIII inhibitors using cyclophosphamide in a patient with systemic lupus erythematosus, 1007

TROTTA F see also Gallerani M

TSUCHIDA T see Matsui K

Тѕикамото Н see Horiuchi Т

TSUTSUMI A see Atsumi T

TUGWELL P see Verhoeven AC

Tumour necrosis factor alpha gene polymorphism, relevance of in rheumatoid arthritis, (L) 681, 682

Tumour necrosis factor alpha, interleukin-4 and soluble CD23 levels. Relationship with CD5 and CD23 expression on B cells in peripheral blood and synovial fluid of rheumatoid arthritis patients, 325

Tumour necrosis factor, assessing role of anti-tumour necrosis factor therapy in the treatment of rheumatoid arthritis. Recapitulation of round-table discussion (S2) 50

TUNACI A see Gül A

TUNACI M see Gül A

TUNCER M see Yetgin S

TURESSON C, JACOBSSON L, BERGSTRÖM U. Extra-articular rheumatoid arthritis: prevalence and mortality, 668

TURNER DE see Woodburn J

Twins and the genetic architecture of osteoarthritis, (E) 583

TVI AVSKY F see Carbone I.

TYNDALL A. Haematological stem cell transplantation in the treatment of severe autoimmune diseases: first experiences from an international project, 774

TZIOUFAS AG see Mavragani CP

UCHIDA S see Manabe N

UHL M see Jacki SH ULLMAN U see Sahly H

Ultrasonic measurement of the thickness of human articular cartilage in situ, 1269

Ultrasonography, the current status of in rheumatology, (E) 195

Ultrasound, preliminary study of aspiration of bone erosion in early rheumatoid arthritis, 329

URBANO-MÁRQUEZ À see Miró O UTHMAN IW see Hajj-Ali RA

UTLEY M see Solymossy C

Uveitis, acute anterior, in a patient with sacroiliitis and acne conglobata, (L) 797

UZIEL Y see Krause I

VAIOPOULOS G see Tatsis G

VALLOW J see Fraser A

VAN BEKKUM DW. Effectiveness and risks of total body irradiation for conditioning in the treatment of autoimmune disease with autologous bone marrow transplantation, 757

VAN DUK H see Lafeber FPJG

VAN DE PUTTE LBA, KROOT EJA, VAN RIEL PLCM. Management of refractory rheumatoid arthritis, (S2) 32

VAN DE PUTTE LBA see also Barreta P VAN DEN BORNE BEE, LANDEWÉ RBM, GOEI THE HS, BREEDVELD FC, Dukmans BAC. Cyclosporin A therapy in rheumatoid arthritis: only strict application of the guidelines for safe use can prevent irreversible renal function loss, 254

VAN DEN BRINK HG see Swaak AJG

VAN DEN WORM E see Lafeber FPJG

VAN DER GRAAFF WL, PRINS APA, NIERS TMH, DIJKMANS BAC, VAN LIER RAW. Quantitation of interferon gamma- and interleukin-4-producing T cells in synovial fluid and peripheral blood of arthritis patients, 214

VAN DER HEIJDE D, BOONEN A, BOERS M, KOSTENSE P, VAN DER LINDEN S. Reading radiographs in chronological order, in pairs or as single films has important implications for the discriminative power of rheumatoid arthritis clinical trials, 1213

VAN DER HEUDE D, DANKERT T, NIEMAN F, RAU R, BOERS M. Reliability and sensitivity to change of a simplification of the Sharp/van der Heijde radiological assessment in rheumatoid arthritis, 941

VAN DER HEIJDEN IM, WILBRINK B, SCHOULS LM, VAN EMBDEN JDA, BREEDVELD FC, TAK PP. Detection of mycobacteria in joint samples from patients with arthritis using a genus-specific polymerase chain reaction and sequence analysis, 547

VAN DER HORST-BRUINSMA IE, HUIZINGA TWJ, LAGAAY EM, HAZES JMW, BREEDVELD FC, SCHREUDER GMTh, TOMSON TAS, ZWINDERMAN AH, VAN ROOD JJ, de VRIES RRP. The influence of a partially HLA-matched blood transfusion on the disease activity of rheumatoid arthritis, 53

VAN DER LELIE H see Slaper-Cortenbach ICM

VAN DER LINDEN MW see Huizinga TWJ VAN DER LINDEN S see van der Heijde D

VAN EMBDEN JDA see van der Heijden IM

VAN GESTEL AM see Albers JMC

VAN HASELEN RA. Research on complementary medicine in rheumatic diseases: the need for better quality studies and reproduction of claimed positive results, (E) 387

VAN HEUGTEN HG see Slaper-Cortenbach ICM

VAN LEEUWEN MA see Posthumus MD

VAN LIER RAW see van der Graaff WL

VAN LOON AM see Blaauw AAM
VAN RIEL PLCM see Albers JMC; Barrera P; Laan RFJM; van de Putte LBA

VAN RUSWUK MH see Posthumus MD

VAN RUSWKUK MH see Brooks P

VAN RINSUM AC see Jacobs JWG

VAN ROOD JJ see van der Horst-Bruinsma

VAN ROON JAG see Lafeber FPJG

VAN ROY JLAM see Lafeber FPJG VAN ROYEN BJ see Bot SDM

VAN SOESBERGEN RM see Drossaers-Bakker KW

VAN VOOREN J-P see Steinfeld SD

VAN 'T HOF MA see Albers JMC

VARON D see Barzilai A

Vascular basis for repetitive strain injury, 636

Vascular involvement. Oxidative stress in systemic lupus erythematosus and allied conditions with, 529

Vasculitic nature of polymyalgia rheumatica, new arguments for, using

positron emission tomography, 444 Vasculitides, virus-associated, (E) 588

Vasculitis

cutaneous necrotizing, erythema nodosum and ankylosing spondylitis, (L) 377

large vessel, simulating Takayasu's arteritis. Sarcoidosis with palpable nodular myositis: periostitis and, (L) 287

necrotizing, secondary to propylthiouracil presenting as purpura fulminans, (L) 790

VEALE D see McGonagle D

VECCHIO P, THOMAS R, HILLS BA. Surfactant treatment for osteoarthritis, (L) 1020

VELICITAT P see Dougados M VENGE P see Torsteinsdóttir I

VERDONCK LF see Slaper-Cortenbach ICM

VERHOEVEN AC, Boers M, Tugwell P. Combination therapy in rheumatoid arthritis, (L) 789

VERSENDAAL J see Kraan MC

VERWEIJ CL, HUIZINGA TWJ. Relevance of tumour necrosis factor alpha gene polymorphism in rheumatoid arthritis, (L) (Reply) 682

VESELY M see Steiner G

VESGA JC see Cuende E

VEYS EM see Dougados M

VIANEN ME see Lafeber FPJG

VIDAL E see Liozon E

VIGNON E see Maillefert JF

VILLARREAL-GARCIA-LOMAS M see Jimenez-Mazuecos J

VILLAVERDE V, BALSA A, CABEZAS JA, FERNÁNDEZ-PRADA M, TORRE A, MOLA EM. Normotensive renal failure in a patient with systemic sclerosis and p-antineutrophil cytoplasmic autoantibodies developed into Paget's disease of bone after immunosuppressive therapy, (L) 190

VILLAVERDE V, MUÑOZ-FERNÁNDEZ S, HIDALGO V, CORTÉS I, FONSECA A, GIJÓN-BAÑOS J, MARTIN-MOLA E. Acute anterior uveitis in a patient with sacriliitis and acne conglobata, (L) 797

VIOLA S see Martini A; Ruperto N

Vitamin D deficiency, acroparasthesia-a typical finding, (L) 482

VLACHOYIANNOPOULOS PG see Swaak AJG

VON KEMPIS J see Jacki SH

Vossen JM, Brinkman DMC, Bakker B, Hoogerbrugge PM, ten Cate R. Rationale for high-dose cyclophosphamide and medium-dose total body irradiation in the conditioning of children with progressive systemic and polyarticular juvenile chronic arthritis before autologous stem cell transplantation, 762

Votsiou A see Tatsis G

WADA M, TATSUO H, BABA H, ASAMOTO K, NOJYO Y. Femoral intercondylar notch measurements in osteoarthritic knees, 554

WAGSTAFF K see Devlin J

WAGURI-NAGAYA Y see Muro H

WAKEFIELD RJ, GIBBON WW, EMERY P. The current status of ultrasonography in rheumatology, (E) 195 WAKEFIELD RJ see also McGonagle D

WALI JP see Handa R

WALKER DJ, KIDD E, HESLOP PS, CHANDLER C. Spontaneous ambulatory activity as a quantifiable outcome measure for rheumatoid arthritis,

WALKER H see O'Duffy EK WALKER JG, LITTLEJOHN GO, MCMURRAY NE, CUTOLO M. Stress system response and rheumatoid arthritis: a multilevel approach, 1050

WALPORT MJ see Naughton MA

WALSH DA. Angiogenesis and arthritis, 103 WALTERS SJ see Brazier JE

WAN J see Carbone L WARD ME see Wilkinson NZ

WARNER T see Brooks P

WATERFIELD JD, McGEER EG, McGEER PL. The peripheral benzodiazepine receptor ligand PK 11195 inhibits arthritis in the MRL-lpr mouse model, 1068

WATSON R see Flanagan N

WEBER M, HASLER P, GERBER H. Sacral insufficiency fractures as an unsuspected cause of low back pain, (L) 90

Weber-Christian disease associated with lupus anticoagulant and anticardiolipin antibodies, (L) 796

WEBLEY M see Persey M WECHSLER B see Généreau T: Papo T

Wegener's granulomatosis

an unusual cause of blindness in, (L) 578 and diabetes insipidus, (L) 684

dilated cardiomyopathy in, (L) 1164

etoposide in, (L) 473

a search for *Pneumocystitis carinii* DNA by polymerase chain reaction on bronchoalveolar lavage fluids from patients with, (L) 1025 small intestinal perforation complicating, (L) 289

splenic infarction in, (L) 1162

WEINBERGER A see Krause I

WEINBLATT ME. The role of current strategies in the future treatment of rheumatoid arthritis, (S2) 19

Weisman MH, Smolen JS. Introduction: future trends in the management of rheumatoid arthritis, (S2) 1

WENDLING D see Maillefert JF; Toussirot E

WESTRA J see Posthumus MD

WHALLETT A see Gillott TJ

WHITE AR, ERNST E. A systematic review of randomized controlled trials of acupuncture for neck pain, 143

WHITE S see Middleton J

WICKI J, OLIVIERI J, PIZZOLATO G, SARASIN F, GUILLEVIN L, DAYER J-M, CHIZZOLINI C. Successful treatment of polyarteritis nodosa related to hepatitis B virus with a combination of lamivudine and interferon alpha, (L) 183

WIEDERKEHR JL see Papo T

Wig N see Handa R

WIJNGAARDEN-DU BOIS see Slaper-Cortenbach ICM

WILBRINK B see van der Heijden IM

WILKINSON ID see Clunic GPR

WILKINSON NZ, KINGSLEY GH, JONES HW, SIEPER J, BRAUN J, WARD ME.

The detection of DNA from a range of bacterial species in the joines
of patients with a variety of arthritides using a nested, broad-range
polymerase reaction chain, 260

WILLIAMS AS see Richards PJ

WILLIAMS B see Emery P WILLIAMS BD see Richards PJ

WILLIAMS P see Solymossy C

WILLIAMSON L, CARTER N. Doctors' handwriting—does size matter?, (L) 683
 WILSON AG, DUFF GW. Relevance of tumour necrosis factor alpha gene polymorphism in rheumatoid arthritis, (L) 681

WILSON H see Denton CP

WINFIELD J see Solymossy C; Squirrell DM

WINTERS JC, SOBEL JS, GROENIER KH, ARENDZEN JH, MEYBOOM-DE JONG B. The long-term course of shoulder complaints: a prospective study in general practice, 160

WITZMANN G see Steiner G

WOBER W. Comparative efficacy and safety of nimesulide and diclofenac in patients with acute shoulder, and a meta-analysis of controlled studies with nimesulide, (S1) 33

WOEHL-KREMER B see Papo T

WOLFE F. Determinants of WOMAC function, pain and stiffness scores: evidence for the role of low back pain, symptom counts, fatigue and depression in osteoarthritis, rheumatoid arthritis and fibromyalgia, 355 WOLFE F see also Maini RN

WOMAC function, pain and stiffness scores, determinants of: evidence for the role of low back pain, symptom counts, fatigue and depression in osteoarthritis, rheumatoid arthritis and fibromyalgia, 355

Wong RWS see Kung AWC

WOODBURN J, TURNER DE, HELLIWELL PS, BARKER S. A preliminary study determining the feasibility of electromagnetic tracking for kinetics at the ankle joint complex, 1260 Work disability in an cohort of patients with seropositive rheumatoid arthritis: a 20 year study, 1138

World Wide Web, insight into rheumatology resources available, (L) 1027, 1028, 1029

WORTHINGTON J see Donn RP

WOTTON SF, DIEPPE PA, DUANCE VC. Type IX collagen immunoreactive peptides in synovial fluids from arthritis patients, 338

WRIGHT I see Pritchard MH

Wrist, histoplasmosis of: a case report, 906

WULFFRAAT NM, Kuis W. Autologous stem cell transplantation: a possible treatment for refractory juvenile chronic arthritis, 764

WULFFRAAT NM, KUIS W, Petty R. Addendum: proposed guidelines for autologous stem cell transplantation in juvenile chronic arthritis, 777 WULFFRAAT NM see also Kuis W; Slaper-Cortenbach ICM

X-ray images, digitalized, Larsen scoring of, 1127

YAMAOKA K see Nakashima H

YANG LY, HUANG WJS, CHEN WP, FU LW, LIN CY. Does parenteral oestrogen therapy flare up disease activity in patients with systemic lupus erythematosus complicated by haemorrhagic cystitis?, (L) 372

YAO JQ, SEEDHOM BB. Ultrasonic measurement of the thickness of human articular cartilage in situ, 1269

YAOUANQ J see Perdriger A

YASUNAGA S see Horiuchi T

YATES DB see Aziz W

YAZICI H see Barnes CG; Ertan S; Hamuryudan V; Korkmaz C

YAZICIOĞLU K see Özgül A

YEBRA-BANGO M see Jimenez-Mazuecos J

YENICESU I see Yetgin S

YETGIN S, OZEN S, SAATCI U, BAKKALOGLU A, TOPALOĞLU R, YENICESU I, OLCAY L, OKUR H, KARAAGAOGLU E, TUNCER M, BESBAS N. Evaluation of tumour necrosis factor alpha, interferon gamma and granulocytemacrophage colony stimulating factor levels in juvenile chronic arthritis, 468

YEUNG SSC see Kung AWC

YIN Z, SIEGERT S, NEURE L, GROLMS M, LIU L, EGGENS U, RADBRUCH A, BRAUN J, SIEPER J. The elevated ratio of interferon gamma-/interleukin-4-positive T cells found in synovial fluid and synovial membrane of rheumatoid arthritis patients can be changed by interleukin-4 but not by interleukin-10 or transforming growth factor beta. 1058

YOSHIDA A see Inokuma S YOUNG A see Solymossy C YOUNG P see Crawford F YURDAKUL S see Hamuryudan V

ZAATARI GS see Hajj-Ali RA

ZAMORANO J see Fernández-Gutiérrez B ZAPHIROPOULOS G see Gillott TJ

ZARDI L see Claudepierre P

ZARDI L see Claudepierre P ZARÓWNY D see Smoleňska Z

ZECCA M see Martini A

ZEIDLER H see Brooks P; Calin A; Dougados M

ZENTILIN P see Filaci G

ZENZ P see Steiner G

ZHANG M see Pelletier J-P

Zidovudine in primary Sjögren's syndrome, 814

ZULIAN F see Ruperto N

ZWINDERMAN AH see van der Horst-Bruinsma



